



Research Article

Epigenetic Insights into Bladder Cancer: The Role of Urine DNA Methylation Assays: Mini Review

Mohamed Hussein

Department of Biomedical Sciences, Dubai Medical College for Girls, Dubai Medical University, United Arab Emirates.

Article History

Submitted: January 02, 2024

Accepted:

Published:

Abstract

Bladder cancer remains a major global health challenge due to its high incidence, recurrence rates, and reliance on invasive diagnostic procedures. While conventional methods like cystoscopy and urine cytology are standard, they often fall short in sensitivity, especially for early-stage disease. Epigenetic alterations, particularly DNA methylation, are now recognized as critical in bladder cancer development and progression. Non-invasive urine-based DNA methylation assays have emerged as promising diagnostic and monitoring tools by detecting tumor-derived molecular changes in exfoliated DNA. Notably, hypermethylation of tumor suppressor gene promoters such as *CDKN2A*, *RASSF1A*, and *DAPK*, as well as hypomethylation in oncogene-associated regions like *MYC* and *CCND1*, have shown strong associations with bladder cancer presence and severity. Recent advancements in detection technologies, including methylation-specific PCR (MSP), droplet digital PCR (ddPCR), and Next-Generation Sequencing (NGS) have significantly improved sensitivity and specificity, enabling earlier diagnosis, better risk stratification, and the development of personalized management strategies. Clinical studies also point to the potential of novel markers like *TWIST1*, *NID2*, and *mDMRTA2* in enhancing diagnostic accuracy. Despite these advances, broader clinical validation and standardized testing protocols are essential for integrating methylation assays into routine clinical practice. These assays represent a transformative shift toward non-invasive, accurate, and patient-centric bladder cancer care.

Keywords

Bladder cancer, DNA methylation, Urine-based diagnostics, Epigenetic biomarkers, Non-invasive assays

Introduction

1.1 Bladder cancer

Bladder cancer is a significant global health concern due to its high incidence, substantial mortality, and the challenges it presents in diagnosis and treatment. It is a heterogeneous disease with varying clinical presentations and outcomes, requiring in-

dividualized management strategies. Men are affected approximately 3 to 4 times more often than women, likely due to differences in exposure to risk factors such as smoking and occupational carcinogens (1). Figure 1. The median age at diagnosis is 73 years, making it a disease predominantly of the elderly. Bladder cancer accounts for an estimated 213,000 deaths annually worldwide. While non-muscle-invasive bladder cancer (NMIBC) has a lower mortality rate, muscle-invasive bladder cancer (MIBC) and metastatic cases contribute significantly to

mortality (2,3).




Figure 1 illustrates bladder cancer staging using the TNM classification, where the "T" component refers to the extent of tumor invasion. Ta indicates a non-invasive papillary tumor confined to the bladder lining, while Tis (carcinoma in situ) refers to a flat, high-grade, non-invasive lesion. T1 tumors invade the connective tissue beneath the lining (lamina propria) but not the muscle. T2 tumors penetrate the muscle layer. T2a involves the inner half, and T2 b the outer half. T3 tumors extend into the peri-vesical (fatty) tissue surrounding the bladder, with T3a indicating microscopic and T3b macroscopic invasion. T4 tumors have spread to nearby organs such as the prostate, uterus, or pelvic wall. This staging helps determine the severity of the disease and guides treatment planning. Created with <https://app.biorender.com/illustrations/683913466895553df7df7bc5>

1.2 The Critical Role of Early Detection in Bladder Cancer Management

Early detection of bladder cancer is pivotal for improving outcomes, as it enables timely interventions that significantly enhance survival rates, prevent disease progression, and reduce the burden of advanced treatments. Early-stage bladder cancer, particularly non-muscle-invasive bladder cancer (NMIBC), has a far better prognosis compared to advanced stages, with a 5-year survival rate of ~77% for localized disease, compared to ~30% for regional spread and <10% for distant metastases. Identifying NMIBC early allows for effective management with intravesical therapies, such as Bacillus Calmette-Guérin (BCG), preventing progression to muscle-invasive bladder cancer (MIBC), which is more aggressive and requires radical treatments (4–6). Early diagnosis broadens treatment options, favoring minimally invasive approaches like transurethral resection of bladder tumors (TURBT) and bladder-preserving therapies, while reducing the need for radical cystectomy and systemic chemotherapy, thereby preserving bladder function and quality of life. It also facilitates better risk stratification, allowing high-risk patients to be closely monitored for recurrence or progression using advanced non-invasive diagnostic tools, such as urine-based DNA methylation assays and NMP22 tests, which improve patient compliance and screening feasibility. Furthermore, early detection minimizes costs associated with lifelong surveillance and advanced disease management and helps prevent metastasis through timely systemic and targeted therapies, ultimately improving survival outcomes and reducing the physical, emotional, and financial burdens on patients (7–10).

1.2.1 Research Methodology

An extensive literature review was carried out using PubMed, Scopus, and Web of Science to identify relevant publications from 2015 to 2025 concerning urine-based DNA methylation assays for bladder cancer. The search utilized keywords such as "bladder cancer," "urine DNA methylation," "non-invasive biomarkers," "epigenetic assays," "qMSP," "ddPCR," and "NGS." Studies were included if they were original research articles or clinical trials involving human participants, focused on the diagnostic or prognostic use of DNA methylation markers in urine, and reported quantitative performance metrics such as sensitivity, specificity, or area under the curve (AUC). Exclusion criteria encompassed studies using non-urine specimens (e.g., tissue or blood), in vitro or animal-only studies, and non-primary research such as reviews. Studies lacking quantitative outcomes related to methylation biomarkers or focusing exclusively on treatment response without diagnostic relevance were also excluded. The initial search identified 299 articles. After screening titles and abstracts, 75 full-text articles were evaluated, resulting in 41 studies that met all inclusion criteria and were analyzed in this review.

1.3 The Role of DNA Methylation in Epigenetic Regulation and Development

DNA methylation is an essential epigenetic mechanism in which a methyl group is added to the 5th carbon of the cytosine ring, primarily within CpG dinucleotides **Figure 2**. This reaction is catalyzed by DNA methyltransferases (DNMTs), including DNMT1 (which maintains methylation during DNA replication) and DNMT3A/3B (which establish new methylation patterns). Methylation in promoter regions typically represses gene transcription by recruiting methyl-CpG-binding domain proteins (MBDs), which compact chromatin and restrict transcriptional access. This process is vital for regulating gene expression, silencing repetitive sequences, and maintaining genomic stability. However, when disrupted, it can contribute to disease development, including cancer (11–14).




Figure 2 illustrates DNA methylation as a heritable epigenetic modification in which a methyl group ($-CH_3$) is covalently added to the 5th carbon position of the cytosine ring, typically within cytosine-guanine (CpG) dinucleotides. The process is catalyzed by DNA methyltransferases (DNMTs), a family of enzymes that play a key role in establishing and maintaining methylation patterns. This modification is essential for regulating gene expression, preserving genomic stability, and guiding cell differentiation and development. Created with <https://app.biorender.com/illustrations/683913466895553df7df7bc5>

DNA methylation plays a vital role in gene regulation by influencing transcriptional activity, genomic stability, and cellular processes. Methylation in gene promoter regions is closely linked to transcriptional repression, as methylated promoters recruit methyl-CpG-binding domain proteins (MBDs) that form complexes with histone-modifying enzymes to compact chromatin, restricting access to the transcriptional machinery. This mechanism is critical during development, such as silencing pluripotency genes in differentiated cells. Additionally, DNA methylation helps maintain genomic stability by silencing repetitive sequences and transposable elements, preventing inappropriate recombination and genomic instability. DNA methylation also ensures genomic stability by silencing repetitive sequences and transposable elements, thereby preventing recombination events and preserving genomic integrity (15). It contributes to X-chromosome inactivation in females, equalizing gene dosage with males, and plays a role in genomic imprinting by regulating parent-specific expression of certain genes essential for normal growth and development. However, aberrant DNA methylation is implicated in various diseases, particularly cancer. Hypermethylation of tumor suppressor gene promoters, such as p16 or MLH1, leads to gene silencing and facilitates uncontrolled cell proliferation. Conversely, global hypomethylation, a hallmark of many cancers, activates oncogenes and transposable elements, increasing genomic instability. Thus, DNA methylation is a dynamic process that maintains cellular homeostasis but, when disrupted, contributes to disease pathogenesis (16–18).

1.4 DNA Methylation and Bladder Cancer: A Dual Role in Genomic Stability and Tumorigenesis

In bladder cancer, DNA methylation patterns become dysregulated, contributing to tumorigenesis and disease progression. Two major types of alterations are commonly observed: global hypomethylation and locus-specific methylation changes (19).

1.4.1 Global Hypomethylation Global hypomethylation involves the loss of methylation in repetitive sequences, transposable elements, and intergenic regions. It is often observed in later stages of bladder cancer. **Genomic Instability:** Hypomethylation activates transposable elements (e.g., LINE-1) and leads to chromosomal instability. **Oncogene Activation:** Hypomethylation in promoter regions of oncogenes results in overexpression, promoting tumor growth and progression. Hypomethylation of LINE-1 sequences is a potential biomarker for bladder cancer prognosis. Hypomethylation in non-coding regulatory regions can alter enhancer activity, leading to abnormal expression of nearby oncogenes or loss of regulatory control (20).

1.4.2 Locus-specific methylation changes involve hypermethylation or hypomethylation of specific genes or genomic regions. These changes are often more directly associated with bladder

cancer development and progression. Hypermethylation occurs predominantly in CpG islands within the promoter regions of tumor suppressor genes. It leads to transcriptional silencing of these genes. Promoter hypermethylation of genes like CDKN2A and RASSF1A is a promising diagnostic and prognostic marker. **Hypomethylation:** Localized hypomethylation can occur in promoter or enhancer regions of oncogenes. It results in the overexpression of genes that drive tumorigenesis. Oncogenes such as MYC and CCND1 can exhibit hypomethylation in their regulatory regions, leading to their upregulation. Hypomethylation in cancer-testis antigens (e.g., MAGE-A1) is frequently observed in bladder cancer (21,22).

1.5 Unlocking the Diagnostic Potential of Urine in Bladder Cancer

Urine serves as an excellent diagnostic medium for bladder cancer due to its accessibility and the non-invasive nature of its collection, making it ideal for repeated sampling and monitoring. As a direct contact medium with the bladder, urine contains a wealth of molecular information shed by tumor cells, allowing for the detection of a variety of biomarkers. These include DNA (e.g., mutations, methylation patterns), RNA (e.g., microRNAs, mRNAs), proteins (e.g., NMP22, cytokeratins), and metabolites, all of which can provide valuable insights into tumor biology and progression. The ease of urine collection reduces patient discomfort compared to invasive procedures like cystoscopy, while its composition reflects dynamic changes in the tumor microenvironment, offering opportunities for early detection, risk stratification, and monitoring of treatment responses. These attributes position urine-based diagnostics as a powerful tool for improving the non-invasive management of bladder cancer (23).

Urine DNA methylation assays have emerged as a promising tool for the non-invasive diagnosis and monitoring of bladder cancer. These assays detect abnormal methylation patterns in cell-free DNA or exfoliated tumor cells present in the urine, offering insights into tumor biology and epigenetic alterations associated with cancer. Aberrant DNA methylation, such as hypermethylation of tumor suppressor gene promoters (e.g., CDKN2A, RASSF1A, DAPK), is a hallmark of bladder cancer and plays a critical role in tumor progression and recurrence. The high sensitivity and specificity of methylation assays make them particularly valuable for detecting bladder cancer at early stages, even before visible tumors are detected by imaging or cystoscopy. Additionally, these assays are ideal for monitoring patients during follow-up to detect residual disease or recurrence, reducing the need for invasive procedures. Advances in technologies such as methylation-specific PCR (MSP), droplet digital PCR (ddPCR), and next-generation sequencing (NGS) have enhanced the reliability and scalability of these tests. Urine DNA methylation assays hold immense potential not only for improving early detection but also for risk stratification and

guiding personalized treatment strategies, making them a key innovation in bladder cancer management (24,25).

Clinical trials have demonstrated the utility of urine-based DNA methylation assays for various clinical objectives in bladder cancer management, including diagnosis, monitoring, and biomarker discovery. For early diagnosis, techniques with high sensitivity, such as quantitative methylation-specific PCR (qMSP) and droplet digital PCR (ddPCR), have been evaluated for their ability to detect low-abundance methylated DNA associated with bladder tumors. For instance, a trial investigating the methylation of CDKN2A and RASSF1A in urine samples reported high sensitivity and specificity for distinguishing bladder cancer patients from healthy controls, even in early-stage disease. These findings highlight the potential of qMSP and ddPCR to facilitate early detection, reducing reliance on invasive cystoscopy and improving patient compliance (25,26).

In the context of disease monitoring, precision techniques like ddPCR and targeted next-generation sequencing (NGS) have shown promise in detecting residual disease and predicting recurrence. A multicenter trial utilizing ddPCR to monitor RASSF10 and APC methylation in urine demonstrated that methylation levels correlate strongly with recurrence risk, providing a non-invasive method for long-term surveillance. Such approaches enable timely intervention, improving outcomes while reducing the need for frequent invasive procedures. Additionally, studies combining ddPCR with other urine biomarkers, such as protein or RNA signatures, have enhanced the predictive power of these assays for recurrence detection. For biomarker discovery, whole-genome or targeted NGS platforms have been pivotal in identifying novel methylation markers for bladder cancer. Trials leveraging NGS to profile urine samples have uncovered multiple methylation candidates, such as ZIC4 and GATA4, that differentiate between bladder cancer subtypes and stages. These biomarkers are being validated in ongoing clinical trials for their diagnostic, prognostic, and predictive utility. Furthermore, NGS-based studies have revealed methylation patterns specific to high-risk bladder cancer, aiding in patient stratification and the development of personalized treatment strategies (23,24).

1.6 Comparative Analysis of Urine DNA Methylation Assays and Other Bladder Cancer Biomarkers

Urine-based DNA methylation assays have emerged as highly promising tools in bladder cancer diagnostics, especially when evaluated alongside established and emerging urinary biomarkers such as cytology, NMP22, UroVysion FISH, and multiplex platforms like Cxbladder. While urine cytology is highly specific (>90%), its low sensitivity, particularly for low-grade tumors, limits its use as a standalone diagnostic (27). NMP22 provides moderate sensitivity (50–70%) but lower specificity and

can yield false positives in benign conditions, whereas UroVysion FISH offers improved accuracy for high-grade tumors with 60–80% sensitivity and 70–85% specificity. Cxbladder, which measures multi-gene mRNA signatures, shows high negative predictive value, making it useful for ruling out malignancy (28). By comparison, DNA methylation assays targeting genes like CDKN2A, RASSF1A, and TWIST1 demonstrate consistently high sensitivity (70–90%) and specificity (85–95%) even for early-stage disease (29). For initial diagnosis in patients with hematuria, combining methylation markers with traditional tests like cytology or NMP22 enhances diagnostic sensitivity without compromising specificity. During follow-up care, these assays also reduce reliance on invasive cystoscopy, with studies confirming their utility in surveillance settings (29, 30). There is also growing interest in combining DNA methylation assays with other biomarker types such as proteins (e.g., NMP22) and RNA signatures (e.g., Cxbladder) to boost accuracy and reduce diagnostic uncertainty. These approaches are particularly beneficial for high-risk patients or when test results are inconclusive, helping to improve clinical decision-making and avoid missed diagnoses or unnecessary procedures (31,32). While molecular tools like UroVysion and Cxbladder offer enhanced performance, they come with higher costs and require specialized infrastructure, limiting their accessibility. In contrast, methylation assays using techniques such as qMSP and ddPCR offer strong diagnostic accuracy with better scalability and cost-efficiency. These assays can be centralized, streamlining laboratory processes and enabling integration with other molecular diagnostics, an advantage for clinics seeking accurate, practical, and affordable bladder cancer testing options (33,34,35).

1.7 CDKN2A, RASSF1A, and Beyond: A Clinical Perspective on Urinary DNA Methylation Testing in Bladder Cancer

Several clinical studies have investigated the potential of urinary DNA methylation biomarkers such as CDKN2A and RASSF1A in the diagnosis of bladder cancer, each contributing valuable insights into assay performance and clinical applicability. A large multicenter study evaluated a methylation-based urine test in over 1,000 patients with hematuria, demonstrating high sensitivity (89.2%) and specificity (87.8%), notably outperforming traditional methods like urine cytology and NMP22 (36). Similarly, a study validated a qMSP-based panel including GHSR and MAL, achieving 80% sensitivity and 93% specificity in urine samples (37). Earlier research reviewed the broader landscape of DNA methylation assays, comparing detection technologies such as ddPCR and NGS, and highlighting challenges related to assay standardization and variability in DNA sources (38). Supporting these findings, another study reported a correlation between CDKN2A and RASSF1A promoter hypermethylation and tumor grade in urothelial carcinoma, reinforcing their prognostic relevance (39). More recently, a novel marker, mDMRTA2, was introduced and validated through a urine-

based assay with strong diagnostic accuracy (40). Collectively, these studies underscore the growing evidence for urinary methylation biomarkers while emphasizing the need for standardized methodologies, larger cohorts, and real-world validation to support clinical adoption.

1.8 Multi-Omics and Machine Learning: Unlocking Precision Oncology in Bladder Cancer

Recent progress in cancer systems biology has underscored the growing impact of multi-omics integration in refining prognostic models for bladder cancer. By merging DNA methylation data with other molecular layers such as transcriptomics and genomic alterations, researchers are gaining a more complete understanding of tumor dynamics. For example, one study used a machine learning-based multi-omics approach to investigate mitotic catastrophe heterogeneity in bladder cancer, identifying distinct molecular subtypes and the gene ANLN as a significant prognostic marker, demonstrating how integrative strategies can uncover meaningful biological patterns (41). Another study combined RNA-seq, miRNA-seq, methylation profiles, and number variations using a transfer learning-based Cox model, resulting in improved prognostic accuracy across various datasets (42). Integrating multi-omics with clinical treatment data has also revealed gene combinations linked to both treatment response and patient outcomes, highlighting new therapeutic opportunities (43). This approach isn't limited to bladder cancer; for instance, a study in endometrial carcinoma identified a novel methylation-based prognostic signature by linking methylation with gene expression data (44). Within bladder cancer specifically, large-scale methylation analyses have identified key prognostic biomarkers (45), and early-stage genomic studies have provided insights into molecular subtypes and potential treatment targets (46).

In parallel, the incorporation of artificial intelligence (AI) and machine learning (ML) into multi-omics analysis has accelerated advances in cancer diagnostics and risk prediction. Deep learning methods have been particularly successful at integrating complex datasets, with a benchmark study revealing that models like moGAT are strong performers in classification, while efmmVAE and efVAE show strength in clustering tasks (47). These findings suggest that selecting the right model depends heavily on the research objective and data structure. Graph-based approaches are also gaining ground—models like LASSO-MOGAT have achieved high accuracy by integrating mRNA, miRNA, and methylation data through Graph Attention Networks and LASSO-based feature selection (48). Together, these developments highlight the promise of AI-driven, multi-omics strategies in advancing precision medicine and improving patient outcomes in oncology.

Conclusions

Bladder cancer continues to pose a significant global health

challenge due to its high prevalence, associated mortality, and dependence on invasive diagnostic techniques. Advances in the understanding of epigenetic mechanisms, particularly DNA methylation, have introduced innovative approaches for diagnosis and monitoring. DNA methylation plays a central role in bladder cancer progression, with global hypomethylation leading to genomic instability and locus-specific methylation changes either silencing tumor suppressor genes or activating oncogenes. These discoveries have facilitated the development of urine-based DNA methylation assays, offering a non-invasive, patient-friendly alternative for early detection, risk assessment, and disease surveillance. Techniques such as methylation-specific PCR (MSP), droplet digital PCR (ddPCR), and next-generation sequencing (NGS) demonstrate high sensitivity and specificity, making them indispensable in clinical and research settings. Clinical studies have highlighted the effectiveness of these assays in identifying early-stage bladder cancer, tracking residual disease, and discovering novel biomarkers for personalized treatment approaches. By minimizing the need for invasive procedures like cystoscopy, urine-based assays improve patient compliance and support more feasible long-term monitoring. Moreover, the incorporation of methylation markers into routine clinical practice holds great potential to revolutionize bladder cancer management by enabling individualized treatment strategies and improving outcomes. Nevertheless, additional research and extensive clinical validation are needed to confirm the scalability, reliability, and cost-effectiveness of these methods in everyday healthcare applications. As the field evolves, urine DNA methylation assays are poised to reshape the future of bladder cancer diagnosis and care, providing a non-invasive, precise, and patient-centered approach.

List of abbreviations

BCG	Bacillus Calmette-Guérin
CpG	Cytosine-phosphate-Guanine
ddPCR	Droplet Digital Polymerase Chain Reaction
DNA	Deoxyribonucleic Acid
DNMT	DNA Methyltransferase
DNMT1	DNA Methyltransferase I
DNMT3A	DNA Methyltransferase 3 Alpha
DNMT3B	DNA Methyltransferase 3 Beta
MBD	Methyl-CpG Binding Domain

MIBC Muscle-Invasive Bladder Cancer

MSP Methylation-Specific Polymerase Chain Reaction

NGS Next-Generation Sequencing

NMIBC Non-Muscle-Invasive Bladder Cancer

NMP22 Nuclear Matrix Protein 22

PCR Polymerase Chain Reaction

qMSP Quantitative Methylation-Specific PCR

RNA Ribonucleic Acid

TURBT Transurethral Resection of Bladder Tumor

Author Contributions

The author conceptualized the study and conducted the literature review. He was responsible for drafting the manuscript, interpreting the data, and revising it critically for important intellectual content. All sections of the manuscript were written and reviewed solely by the author. The author approved the final version of the manuscript and agrees to be accountable for all aspects of the work.

Availability of Data and Materials

The data supporting the findings of this study are available from the corresponding author upon reasonable request.

Conflict of Interest

No Conflict of Interest

Funding

This research received no external funding.

Acknowledgments

The author would like to thank Dubai Medical College for girls for their support during the study.

The author used [BioRender software](#) for creation of figures.

References

1. Dyrskjot L, Hansel DE, Efstathiou JA, Knowles MA, Galsky MD, Teoh J, et al. Bladder cancer. *Nat Rev Dis Primers*. 2023 Oct 26;9(1):58.
2. Lidagoster S, Ben-David R, De Leon B, Sfakianos JP. BCG and Alternative Therapies to BCG Therapy for Non-Muscle-Invasive Bladder Cancer. *Current Oncology*. 2024 Feb 16;31(2):1063–78.
3. Xu Y, Luo C, Wang J, Chen L, Chen J, Chen T, et al. Application of nanotechnology in the diagnosis and treatment of bladder cancer. *J Nanobiotechnol*. 2021 Dec;19(1):393.
4. Siracusano S, Rizzetto R, Porcaro AB. Bladder cancer genomics. *Urologia*. 2020 May;87(2):49–56.
5. Dobruch J, Oszczudłowski M. Bladder Cancer: Current Challenges and Future Directions. *Medicina*. 2021 Jul 24;57(8):749.
6. Hamad J, McCloskey H, Milowsky MI, Royce T, Smith A. Bladder preservation in muscle-invasive bladder cancer: a comprehensive review. *Int braz j urol*. 2020 Mar;46(2):169–84.
7. Holzbeierlein JM, Bixler BR, Buckley DI, Chang SS, Holmes R, James AC, et al. Diagnosis and Treatment of Non-Muscle Invasive Bladder Cancer: AUA/SUO Guideline: 2024 Amendment. *J Urol*. 2024 Apr;211(4):533–8.
8. Babjuk M, Burger M, Capoun O, Cohen D, Compérat EM, Dominguez Escrig JL, et al. European Association of Urology Guidelines on Non-muscle-invasive Bladder Cancer (Ta, T1, and Carcinoma in Situ). *European Urology*. 2022 Jan;81(1):75–94.
9. Khansary S, Tavilani H, Ghasemi H. Gender, Bladder Cancer Healthcare and Burden of COVID-19. *Cancer Investigation*. 2023 Jan 2;41(1):58–69.
10. Tan WS, Steinberg G, Witjes JA, Li R, Shariat SF, Roupret M, et al. Intermediate-risk Non-muscle-invasive Bladder Cancer: Updated Consensus Definition and Management Recommendations from the International Bladder Cancer Group. *European Urology Oncology*. 2022 Oct;5(5):505–16.
11. Koch A, Joosten SC, Feng Z, De Ruijter TC, Draht MX, Melotte V, et al. Analysis of DNA methylation in cancer: location revisited. *Nat Rev Clin Oncol*. 2018 Jul;15(7):459–66.
12. Kulis M, Esteller M. DNA Methylation and Cancer. In: *Advances in Genetics* [Internet]. Elsevier; 2010 [cited 2024 Nov 18]. p. 27–56. Available from: <https://linkinghub.elsevier.com/retrieve/pii/B9780123808660600022>
13. Moore LD, Le T, Fan G. DNA Methylation and Its Basic Function. *Neuropsychopharmacol*. 2013 Jan;38(1):23–38.
14. Law PP, Holland ML. DNA methylation at the crossroads of gene and environment interactions. Blewitt M, editor. *Essays in Biochemistry*. 2019 Dec 20;63(6):717–26.
15. Martisova A, Holcakova J, Izadi N, Sebuyoya R, Hrstka R, Bartosik M. DNA Methylation in Solid Tumors: Functions and Methods of Detection. *IJMS*. 2021 Apr 19;22(8):4247.

16. Ortega-Recalde O, Hore TA. DNA methylation in the vertebrate germline: balancing memory and erasure. Blewitt M, editor. *Essays in Biochemistry*. 2019 Dec 20;63(6):649–61.
17. Mattei AL, Bailly N, Meissner A. DNA methylation: a historical perspective. *Trends in Genetics*. 2022 Jul;38(7):676–707.
18. Nishiyama A, Nakanishi M. Navigating the DNA methylation landscape of cancer. *Trends in Genetics*. 2021 Nov;37(11):1012–27.
19. Wu J, Lin Y, Yang K, Liu X, Wang H, Yu T, et al. Clinical effectiveness of a multitarget urine DNA test for urothelial carcinoma detection: a double-blinded, multicenter, prospective trial. *Mol Cancer*. 2024 Mar 19;23(1):57.
20. Wever BMM, Bach S, Tibbesma M, Ter Braak TJ, Wajon D, Dickhoff C, et al. Detection of non-metastatic non-small-cell lung cancer in urine by methylation-specific PCR analysis: A feasibility study. *Lung Cancer*. 2022 Aug;170:156–64.
21. Guo RQ, Xiong GY, Yang KW, Zhang L, He SM, Gong YQ, et al. Detection of urothelial carcinoma, upper tract urothelial carcinoma, bladder carcinoma, and urothelial carcinoma with gross hematuria using selected urine-DNA methylation biomarkers: A prospective, single-center study. *Urologic Oncology: Seminars and Original Investigations*. 2018 Jul;36(7):342.e15-342.e23.
22. Vener T, Derecho C, Baden J, Wang H, Rajpurohit Y, Skelton J, et al. Development of a Multiplexed Urine Assay for Prostate Cancer Diagnosis. *Clinical Chemistry*. 2008 May 1;54(5):874–82.
23. Fux R, Kloor D, Hermes M, Röck T, Proksch B, Grenz A, et al. Effect of acute hyperhomocysteinemia on methylation potential of erythrocytes and on DNA methylation of lymphocytes in healthy male volunteers. *American Journal of Physiology-Renal Physiology*. 2005 Oct;289(4):F786–92.
24. Roobol MJ, Bangma CH, El Bouazzaoui S, Franken-Raab CG, Zwarthoff EC. Feasibility study of screening for bladder cancer with urinary molecular markers (the BLU-P project). *Urologic Oncology: Seminars and Original Investigations*. 2010 Nov;28(6):686–90.
25. Renard I, Joniau S, Van Cleynenbreugel B, Collette C, Naômé C, Vlassenbroeck I, et al. Identification and Validation of the Methylated TWIST1 and NID2 Genes through Real-Time Methylation-Specific Polymerase Chain Reaction Assays for the Noninvasive Detection of Primary Bladder Cancer in Urine Samples. *European Urology*. 2010 Jul;58(1):96–104.
26. Witjes JA, Morote J, Cornel EB, Gakis G, Van Valenberg FJP, Lozano F, et al. Performance of the Bladder EpiCheck™ Methylation Test for Patients Under Surveillance for Non-muscle-invasive Bladder Cancer: Results of a Multicenter, Prospective, Blinded Clinical Trial. *European Urology Oncology*. 2018 Sep;1(4):307–13.
27. Oyaert M, Van Praet C, Delrue C, Speckaert MM. Novel Urinary Biomarkers for the Detection of Bladder Cancer. *Cancers (Basel)*. 2025 Apr 10;17(8):1283. doi: 10.3390/cancers17081283. PMID: 40282460; PMCID: PMC12025552.
28. Magee D, Tharakan N, Yuiminaga Y. Validation of Cxbladder® Triage and Monitor as an Adjunct to Urothelial Carcinoma Diagnosis and Surveillance in a Single Centre. *Res Rep Urol*. 2025 Mar 18;17:87-94. doi: 10.2147/RRU.S516994. PMID: 40129475; PMCID: PMC11930626.
29. Jeong IG, Yun SC, Ha HK, Kang SG, Lee S, Park S, Sung HH, Kim SI, Hwang EC, Moon KC, Kwak C. Urinary DNA Methylation Test for Bladder Cancer Diagnosis. *JAMA Oncol*. 2025 Mar 1;11(3):293-299. doi: 10.1001/jamaoncol.2024.6160. PMID: 39883469; PMCID: PMC11783243.
30. Yao Z, Wang T, Liu J, Zhou Z, Zhang Y. Diagnostic accuracy of cytology and urine methylation test in patients with non-muscle invasive bladder cancer: a systematic review and meta-analysis. *Front Oncol*. 2024 Sep 3;14:1412346. doi: 10.3389/fonc.2024.1412346. PMID: 39290246; PMCID: PMC11405155.
31. Witjes JA, Morote J, Cornel EB, Gakis G, van Valenberg FJP, Lozano F, Sternberg IA, Willemsen E, Hegemann ML, Paitan Y, Leibovitch I. Performance of the Bladder EpiCheck™ Methylation Test for Patients Under Surveillance for Non-muscle-invasive Bladder Cancer: Results of a Multicenter, Prospective, Blinded Clinical Trial. *Eur Urol Oncol*. 2018 Sep;1(4):307-313. doi: 10.1016/j.euo.2018.06.011. Epub 2018 Jul 17. PMID: 31100252.
32. Renard I, Joniau S, van Cleynenbreugel B, Collette C, Naômé C, Vlassenbroeck I, Nicolas H, de Leval J, Straub J, Van Criekeing W, Hamida W, Hellel M, Thomas A, de Leval L, Bierau K, Waltregny D. Identification and validation of the methylated TWIST1 and NID2 genes through real-time methylation-specific polymerase chain reaction assays for the noninvasive detection of primary bladder cancer in urine samples. *Eur Urol*. 2010 Jul;58(1):96-104. doi: 10.1016/j.eururo.2009.07.041. Epub 2009 Aug 5. PMID: 19674832.
33. Beijert IJ, Wever BMM, Hentschel AE, van den Burgt Y, Kauer PC, Lissenberg-Witte BI, van Moorselaar RJA, Steenbergen RDM, Nieuwenhuijzen JA. Bladder cancer detection in urine by novel methylation markers. *Sci Rep*. 2024 Nov 20;14(1):28705. doi: 10.1038/s41598-024-77781-0. PMID: 39567591; PMCID: PMC11579363.

34. Roobol MJ, Bangma CH, el Bouazzaoui S, Franken-Raab CG, Zwarthoff EC. Feasibility study of screening for bladder cancer with urinary molecular markers (the BLU-P project). *Urol Oncol*. 2010 Nov-Dec;28(6):686-90. doi: 10.1016/j.urolonc.2009.12.002. PMID: 21062653.
35. Thielemans R, Speeckaert R, Delrue C, De Bruyne S, Oyaert M, Speeckaert MM. Unveiling the Hidden Power of Uromodulin: A Promising Potential Biomarker for Kidney Diseases. *Diagnostics (Basel)*. 2023 Sep 28;13(19):3077. doi: 10.3390/diagnostics13193077. PMID: 37835820; PMCID: PMC10572911.
36. Jeong IG, Yun SC, Ha HK, Kang SG, Lee S, Park S, Sung HH, Kim SI, Hwang EC, Moon KC, Kwak C. Urinary DNA Methylation Test for Bladder Cancer Diagnosis. *JAMA Oncol*. 2025 Mar 1;11(3):293-299. doi: 10.1001/jamaoncol.2024.6160. PMID: 39883469; PMCID: PMC11783243.
37. Hentschel AE, Beijert IJ, Bosschieter J, Kauer PC, Vis AN, Lissenberg-Witte BI, van Moorselaar RJA, Steenberg RDM, Nieuwenhuijzen JA. Bladder cancer detection in urine using DNA methylation markers: a technical and prospective preclinical validation. *Clin Epigenetics*. 2022 Feb 5;14(1):19. doi: 10.1186/s13148-022-01240-8. PMID: 35123558; PMCID: PMC8818199.
38. Larsen LK, Lind GE, Guldberg P, Dahl C. DNA-Methylation-Based Detection of Urological Cancer in Urine: Overview of Biomarkers and Considerations on Biomarker Design, Source of DNA, and Detection Technologies. *Int J Mol Sci*. 2019 May 30;20(11):2657. doi: 10.3390/ijms20112657. PMID: 31151158; PMCID: PMC6600406.
39. Bilgrami SM, Qureshi SA, Pervez S, Abbas F. Promoter hypermethylation of tumor suppressor genes correlates with tumor grade and invasiveness in patients with urothelial bladder cancer. *Springerplus*. 2014 Apr 5;3:178. doi: 10.1186/2193-1801-3-178. PMID: 24790823; PMCID: PMC4000596.
40. Deng L, Chao H, Deng H, Yu Z, Zhao R, Huang L, Gong Y, Zhu Y, Wang Q, Li F, Liu L, He L, Tang Z, Liao C, Qi Y, Wang X, Zeng T, Zou H. A novel and sensitive DNA methylation marker for the urine-based liquid biopsies to detect bladder cancer. *BMC Cancer*. 2022 May 6;22(1):510. doi: 10.1186/s12885-022-09616-y. PMID: 35524222; PMCID: PMC9077853.
41. Zhang J, Chen J, Xu M, et al. Exploring prognostic DNA methylation genes in bladder cancer: a comprehensive analysis. *Discov Onc*. 2024;15:331. <https://doi.org/10.1007/s12672-024-01206-7>
42. Prip F, Lamy P, Lindskrog SV, et al. Comprehensive genomic characterization of early-stage bladder cancer. *Nat Genet*. 2025;57:115-125. <https://doi.org/10.1038/s41588-024-02030-z>
43. Xu Y, Sun X, Liu G, Li H, Yu M, Zhu Y. Integration of multi-omics and clinical treatment data reveals bladder cancer therapeutic vulnerability gene combinations and prognostic risks. *Front Immunol*. 2024;14:1301157. doi:10.3389/fimmu.2023.1301157
44. Cao L, Ma X, Rong P, et al. Comprehensive analysis of DNA methylation and transcriptome to identify PD-1-negative prognostic methylated signature in endometrial carcinoma. *Dis Markers*. 2022;2022:3085289. doi:10.1155/2022/3085289
45. Jeong IG, Yun SC, Ha HK, et al. Urinary DNA methylation test for bladder cancer diagnosis. *JAMA Oncol*. 2025;11(3):293-299. doi:10.1001/jamaoncol.2024.6160
46. Hentschel AE, Beijert IJ, Bosschieter J, et al. Bladder cancer detection in urine using DNA methylation markers: a technical and prospective preclinical validation. *Clin Epigenetics*. 2022;14(1):19. doi:10.1186/s13148-022-01240-8
47. Leng D, Zheng L, Wen Y, et al. A benchmark study of deep learning-based multi-omics data fusion methods for cancer. *Genome Biol*. 2022;23:171. <https://doi.org/10.1186/s13059-022-02739-2>
48. Alharbi F, et al. LASSO-MOGAT: a multi-omics graph attention framework for cancer classification. *Acad Biol*. 2024;2(3). doi:10.20935/AcadBiol7325