

Disclaimer: This is not the final version of the article. Changes may occur when the manuscript is published in its final format.

Unusual arrangement of catalytic loops of the alcoholdehydrogenase enzyme during the adsorption process on a graphitic carbon surface

Kholmirzo T. Kholmurodov^{1,2,3,4,*}, Ivan A. Baigunov², Pavel P. Gladyshev²,
Mirzoaziz A. Husenzoda⁵, Hanan G. Elhaes⁶, Medhat A. Ibrahim⁷

¹*Frank Laboratory of Neutron Physics, Joint Institute for Nuclear Research, 141980, Dubna, Moscow Region, Russia*

²*Department of Chemistry, New Technologies and Materials, Dubna State University, 141980, Dubna, Moscow Region, Russia*

³*Department of Fundamental Nuclear Interactions, Faculty of Physics, Lomonosov Moscow State University, 119991, Moscow, Russia*

⁴*S.U.Umarov Physical-Technical Institute (PhTI), Aini ave. 299/1, Dushanbe, 734063, Republic of Tajikistan*

⁵*Department of Technical Operation of Air Transport, Faculty of Transport and Road Infrastructure, Tajik Technical University, Academicians Radjabov str., 10, Dushanbe Republic of Tajikistan, 734042*

⁶*Physics Department, Faculty of Women for Arts, Science and Education, Ain Shams University, Cairo, 11757, Egypt*

⁷*Molecular Spectroscopy and Modeling Unit, Spectroscopy Department, National Research Centre, 35 El-Behouth St., Dokki, Giza, 12622, Egypt*

*Corresponding author. Email: kholmirzo@gmail.com

Kholmirzo Kholmurodov <https://orcid.org/0000-0002-9415-8276>

Ivan Baigunov <https://orcid.org/0009-0000-8380-6218>

Pavel Gladyshev <https://orcid.org/0000-0002-7449-4475>

Hanan Elhaes <https://orcid.org/0000-0002-1234-5431>

Medhat IBRAHIM <https://orcid.org/0000-0002-9698-0837>

Mirzoaziz Khusenov <https://orcid.org/0000-0003-1220-2139>

Disclaimer: This is not the final version of the article. Changes may occur when the manuscript is published in its final format.

ABSTRACT

Molecular dynamics simulations (MDs) were conducted to investigate the alcohol dehydrogenase (ADH) enzyme and its cofactor, nicotinamide adenine dinucleotide (NAD), solvated in an aqueous system on a graphite-carbon surface. The study focuses on key structural aspects of ADH, particularly its conformational changes, including the rotation of the catalytic domain, the coenzyme binding domain, and the rearrangement of the active center required for catalytic activation. MDs were performed for 100 ns to track the conformational and rotational changes of the ADH+NAD complex in an aqueous environment. The diffusion process of ADH+NAD was observed for approximately 50–60 ns, after which the complex made contact with the graphite surface, leading to final relaxation. Upon adsorption of ADH onto the carbon surface, NAD exhibited fluctuations between 55 and 60 ns, coinciding with the opening of two catalytic loops associated with the NAD-binding region. Experimental and kinetic studies suggest that these structural changes may significantly impact the enzymatic activity of ADH in the presence of a carbon surface.

Keywords: proteins' conformation, water solution, graphitic carbon surface, alcohol dehydrogenase enzyme, molecular dynamics, adsorption processes

Disclaimer: This is not the final version of the article. Changes may occur when the manuscript is published in its final format.

INTRODUCTION

The enzyme alcohol dehydrogenase (ADH) of yeast is one of the interesting objects in the chemical, biological, pharmaceutical, electrochemical, agrochemical and aromatic industries [1-5]. For example, in [5], a successful preparation of hexanal catalyzed by ADH from baker's yeast was carried out. Essential to the functionality of ADH is that it must be added in stoichiometric amounts and cannot be replaced by more economical synthetic products. At the same time, the coenzyme NAD⁺ (nicotinamide adenine dinucleotide) is widely used in biocatalytic oxidations catalyzed by ADH. The most intriguing in the structural aspect of ADH, as shown by X-ray crystallography, is that ADH undergoes global conformational changes (it does not matter when binding the NAD⁺ – oxidized form or the NADH – reduced form of NAD), including rotation of the catalytic domain relative to the coenzyme binding domain and rearrangement of the active center to form a catalytically active enzyme. A change in conformation requires the presence of a complete coenzyme and depends on various chemical or mutational substitutions, which can increase catalytic activity by changing the kinetics of isomerization and the rate of dissociation of coenzymes [2-6].

As for the structural aspect of the enzyme, the orientation of ADH on various sorbents and electrically conductive matrices was studied using experimental observations and mathematical modeling of the protein dependence on the pH of the solution. As is known, the deactivation of the enzyme is caused by unsuitable conditions (temperature, pH), which leads to a change in the activity of the enzyme ADH. In relation to the problems of modeling the effect of pH on the orientation of protein sorption, the sorption behavior of the ADH enzyme based on specific interactions of protein groups with groups on the surface and its charge was previously studied in [7-15]. It was assumed that the conformation of the ADH enzyme corresponds to its conformation in the crystal structure and does not change when the surrounding solution and the surface of the sorbent change, which is a rather rough approximation. The implementation of various options includes, for example, the immobilization of a two-substrate enzyme on the surface of electrode materials [7-15].

However, it should be noted that the experimental study of the above issues is difficult. Therefore, in recent years, computational and simulation analysis methods have been widely used for these purposes [16-23]. In this work, we used computer molecular dynamics (MD) modeling to study the structural conformational changes of the enzyme ADH and its cofactor NAD, which occur in aqueous solution when interacting with a graphite carbon surface. The MD analysis data provide

Disclaimer: This is not the final version of the article. Changes may occur when the manuscript is published in its final format.

a significant extension of the initial basic model, thereby allowing the protein conformation to be changed in the region of titrated amino acid residues of ADH in atomic/molecular details. The identification of the characteristic conformation of key titrated amino acids may become a necessary stage for further research and the implementation of a numerical experiment, which will be carried out by varying the pH and surface charge values [2, 7-15].

Of course, our current research correlates with several parallel theoretical and experimental works concerning the immobilization and adsorption of enzymes on a carbon surface, the location and fixation of enzymes on carbon platforms, as well as on the surfaces of bioanodes, etc. It is worth noting that the choice of carbon surfaces and platforms is due to good control of the location of enzymes on the surface with very low enzyme consumption. To date, the kinetic behavior of the ADH enzyme in a solution fixed on carbon platforms has been studied; However, there are few comparative studies of the ADH enzyme that take into account the kinetic behavior of the enzyme, despite the studies of the structure and kinetic mechanism of immobilized enzymes available in the literature. Kinetic parameters obtained for the ADH enzyme attached to a carbon surface (carbon fiber paper) indicate a decrease in activity after the immobilization and fixation process, a significant loss of enzymatic activity observed after immobilization, even while maintaining the affinity between enzymes and their substrates and coenzymes. We believe that MD observations of the structural behavior of ADH+NAD, diffusion and adsorption processes correlate well with the kinetic parameters obtained for the enzyme fixed with ADH on a carbon platform with various kinetic parameters (mainly the concentration of substrates and coenzyme), which indicates a change (loss or decrease) in enzymatic activity fixed enzymes after the immobilization process on the surface. In particular, the observations obtained in our current study may correlate well with practical applications, for example, with the growing interest of biotechnologists in the use of immobilized enzymes such as ADH for many purposes, bioremediation, sensors and biofuel elements, etc.

Materials and Methods

MOLECULAR DYNAMICS SIMULATIONS


In this section, we have implemented the basic MD-AMBER production simulation (CPU/GPU) (also common with many other types of simulation based on processors and GPUs to perform MD simulation with AMBER). We used molecular dynamics (MD) modeling with the

Disclaimer: This is not the final version of the article. Changes may occur when the manuscript is published in its final format.

Amber 18 code (CPU/GPU environment) for the PDB ID: 3COS of the crystal structure of human alcohol dehydrogenase Class II [24]. A detailed description of the main parameters and algorithms used in computer modeling of molecular dynamics is given in our previously published article in [25]. For the ADH+NAD+water+graphite carbon surface system, the general molecular structure shown below in Fig. 1(a-b) was obtained almost manually, step by step. And our simulated system presented no particular difficulties in preparation stage, without using special software or a database of force fields beyond the MD-AMBER computing environment. Thus, to keep the surface atom positions fixed we have used [25] the restraint options (`ntr=1`, `restraint_wt=25`, `restraintmask=':383-35140'`), so far the positional restraint was varied from 5 to 25 kcal mol⁻¹ Å⁻² for the several set of the MD calculations. It's worth noting that the `ntr=1` means that we have turned on position restraints and therefore have to specify via GROUP input the atoms (graphitic carbon surfaces) were restrained as well as the force constant. In the MD run, after specification of the namelists, a title is given, followed by the force constant for the restraint (in kcal mol⁻¹ Å⁻²) and then a specification of surface carbon (C) residues or atoms to restrain. Residues can be specified using the "RES" keyword. We have chosen a force constant of 25 kcal mol⁻¹ Å⁻² and restrained the surrounding ADH+NAD + water confined C-surfaces residues were through 383-35140 (from 1 to 382 the correspondent numerations stand for the ADH+NAD enzyme; a total number of atoms for the ADH+NAD + water + carbon surfaces was 157371). It's worth also noting that the ADH is a Zn(II)-bound metalloenzyme, and during the adsorption process in ADH the position of the metal bound to Zn(II) was fixed, as in the original PDB file (ID: 3COS of the crystal structure of human alcohol dehydrogenase class II ADH). It was necessary to maintain the position of Zn(II) inside the enzyme pocket of ADH+NAD, thereby restraining the relative position of the metal bound to Zn(II) during the entire adsorption process. So far, the position of the Zn(II)-bound metal has not been violated, since the main attention has been paid to the orientation sorption of the ADH+NAD and the behavior of the NAD co-enzyme due to the conformational behavior of the catalytic loops of the enzyme relaxed equilibrated structure.

The MD modeling on the molecular system of ADH+NAD+water+carbon surface, equilibrated at a target temperature of 303 K, was carried out in three stages [25]: energy minimization, NVT and NPT relaxation procedures (Fig. 1)

Disclaimer: This is not the final version of the article. Changes may occur when the manuscript is published in its final format.



Figs. 1(a-b). The general molecular design of the ADH+NAD+water+carbon graphite surface system.

Disclaimer: This is not the final version of the article. Changes may occur when the manuscript is published in its final format.

RESULTS AND DISCUSSION

Fig. 2 shows the resulting pictures of the adsorption of ADH+NAD on the surface of graphite carbon during prolonged 100 ns of dynamic changes from (a) the initially relaxed state to (b–c) intermediate states and (d) the final equilibrium condition.




Fig. 2 (a-d). Absorption of ADH+NAD on carbon graphite (C-surface) for 100 ns dynamical and conformational changes. The upper images show the position of ADH+NAD before adsorption on carbon surfaces (about 60 ns); the lower images show the approach of

Disclaimer: This is not the final version of the article. Changes may occur when the manuscript is published in its final format.

ADH+NAD and the close contact formation with the C-surface. Two catalytic ADH loops are enlarged in size.


The enzyme ADH+NAD undergo multimillion conformational and rotational changes before adsorption on this graphite carbon (C-surface) in order to finally capture and relax on the surface. The dynamics of ADH+NAD adsorption on the graphite carbon surface was monitored using MD/AMBER calculations and Visual Molecular Dynamics (VMD) software. For each MD model, we performed calculations in 100 ns using the module "*pmemd.cuda*". One of the nontrivial events of the conformational structural dynamics of the entire ADH+NAD+water system / C-surface and tracking of individual amino acid residues should be the behavior of the catalytic loops of the enzyme. Fig. 2 show the dynamic patterns of the ADH+NAD / C-surface and the adsorption processes, accompanied by a gradual change in the orientation of the two catalytic loops of the ADH+NAD molecule relative to the graphite surface. An important observation is that these two catalytic loops are located close inside the ADH+NAD molecule, and when we reach the adsorbing graphite surface, we can see the separation of these loops from each other. A visual description of the entire process has shown in Fig. 2, where two enlarged loops separates from each other and, consequently, open important catalytic ADH fragments as a result of the influence of the adsorbing C-surface of these two enzyme loops.

Further, Figs. 3 show the location of the above-mentioned two catalytic loops of the ADH+NAD molecule relative to the graphite surface. Two terminal amino acid residues THR290 and ILE317 are highlighted here, where Δ is introduced, the distance between these two terminal amino acids.


Fig. 3(a) shows the location of the catalytic loops during adsorption of the ADH+NAD molecule on the carbon surface of graphite in the final (100 ns) state. Fig. 3(b) shows the results of calculations of the dynamics of the distance between the catalytic loops of ADH+NAD on the carbon surface of graphite as a function of time. The results in Fig. 3 clearly confirm the important observation mentioned above that the two catalytic loops separate from each other when they reach the adsorbing surface of graphite, whereas initially they were located close to each other inside the ADH+NAD molecule.

Disclaimer: This is not the final version of the article. Changes may occur when the manuscript is published in its final format.

(a)



(b)




Figs. 3(a-b). The positions of the ADH+NAD catalytic loops on the carbon graphite surface (a). Dynamics of the distance between the ADH+NAD catalytic loops as a function of time (b).

The above observation and the presented data on the dynamics of the catalytic loops of ADH+NAD on the carbon surface correlate with the dynamical changes and rotations of the

Disclaimer: This is not the final version of the article. Changes may occur when the manuscript is published in its final format.

coenzyme NAD inside the ADH molecule. Using the data in Fig. 3, we also estimated the conformational changes of NAD during adsorption of ADH on the surface and the open gap between the catalytic loops. In Fig. 4, we noted the positions of the atoms at the beginning and end of the chain NAD, as well as the atoms in the central region NAD.



Figs. 4(a-b). The structure of NAD with the arrangement of atoms from the beginning to the end of the molecular chain (a). The dynamics of the angle of the structure of NAD as a function of time (b).

Fig. 4 demonstrate a method for estimating the angle of γ - vibration and rotation of coal in the three-dimensional structure of the NAD molecule, where auxiliary vectors are introduced and the starting points are designated $A \sim (PA, O3, PN)$, $B \sim N_{begin}$, $C \sim N_{end}$ with the results of calculating the dynamical changes in the angle of γ of the NAD molecule depending on time. They are shown in Fig. 4(b) below. It can be seen that the dynamical changes of γ with time are completely correlated with the dynamics of the change in the distance between the catalytic loops of ADH +

Disclaimer: This is not the final version of the article. Changes may occur when the manuscript is published in its final format.

NAD on the carbon surface of graphite. The comparison of the results in Fig. 3 and Figs. 4 is straightforward. Thus, these results are the key features of the entire process of adsorption of ADH+NAD / C-surface, which is accompanied by a non-trivial structural transformation of the coenzyme NAD, correlating with the behavior of the catalytic loops of the enzyme ADH.

Further, below in Fig. 5, the orientations of the three titrated amino acid residues ASP54, HIS252 and ASP268 with time are shown.




Fig. 5. Dynamics of orientation of two catalytic loops and three titrated amino acid residues ASP54, HIS252 and ASP268 over time.


It is worth noting that for ADH+NAD+water+graphite surface, as described above, all calculations were performed at pH =7 in an aqueous medium. The experimental data and analysis of the kinetic behavior of the ADH enzyme in solution as a function of pH and temperature, have shown that the highest activity is achieved in the pH range 7.0–8.0 and an optimal temperature about 303 K [4-7, 26-28]. At pH 7, the protonation states of the catalytic triad, histidine (HIS), aspartate (ASP), and glutamate (GLU), are determined by their respective pKa values. Histidine has an imidazole side chain with a pKa of approximately 6.0, meaning that at physiological pH, it exists primarily in its deprotonated, neutral form. Aspartate (ASP) and glutamate (GLU), with pKa values of around 3.9 and 4.2, respectively, are both fully deprotonated at pH 7.

In Fig. 6, we calculated the dynamics of the angle of γ – internal vibration and rotation (N-CA-OD2) of three titrated amino acid residues ASP54 (a), HIS252 (b) and ASP268 (c). As in the


Disclaimer: This is not the final version of the article. Changes may occur when the manuscript is published in its final format.

case of the molecule above, we introduced auxiliary vectors, thereby thus, choosing the places of evaluation of amino acid residues in the form of: A~(CA) – the middle atom of the residue; B~N_{begin} (NH3-end) – the residue of one terminal atom; C~OD2 (NE2) – the residue of another terminal atom. From the results shown in Fig. 6 it can be seen that the dynamic behavior and the change in γ over time fully correlate with the dynamics of the change in the distance between the catalytic loops of ADH + NAD on the carbon surface of graphite, as shown in Fig. 3(b), as well as in 4(b).


(a)



(b)




(c)



Disclaimer: This is not the final version of the article. Changes may occur when the manuscript is published in its final format.

Figs. 6(a-b). Dynamics of the angle of γ – internal vibration and rotation (N-CA-OD2) of three titrated amino acid residues ASP54 (a), HIS252 (b) and ASP268 (c) in time.

Figs. 7 show the location of the catalytic loops during adsorption of the ADH+NAD molecule on the carbon surface of graphite in the initial (a) state, (b) in the intermediate (50 ns) and (c) in the final (100 ns) states.



Figs. 7(a-c). The positions of the ADH+NAD catalytic loops on the graphite carbon surface in the initial (a), intermediate (50 ns) (b) and final (100 ns) states (c).

Disclaimer: This is not the final version of the article. Changes may occur when the manuscript is published in its final format.

Fig. 8 demonstrates the procedure for calculating the estimate of the angle of γ - vibration and rotation of coal in the three-dimensional structure of the molecule NAD, where auxiliary vectors are introduced and the starting points are designated A~(PA,O3,PN), B~N_{begin}, C~N_{end}. The dynamical changes in the angle γ of the NAD molecule as a function of time correlate with the dynamics of changes in the distance between the catalytic loops of ADH+NAD on the carbon surface of graphite:

$$\cos \gamma = \frac{(\overline{AB}, \overline{AC})}{|\overline{AB}| \cdot |\overline{AC}|}$$

$$\overline{AB} = (x_B - x_A, y_B - y_A, z_B - z_A), \quad \overline{AC} = (x_C - x_A, y_C - y_A, z_C - z_A)$$

$$(\overline{AB}, \overline{AC}) = (x_B - x_A) \cdot (x_C - x_A) + (y_B - y_A) \cdot (y_C - y_A) + (z_B - z_A) \cdot (z_C - z_A)$$

$$|\overline{AB}| = \sqrt{(x_B - x_A)^2 + (y_B - y_A)^2 + (z_B - z_A)^2}$$

$$|\overline{AC}| = \sqrt{(x_C - x_A)^2 + (y_C - y_A)^2 + (z_C - z_A)^2}$$

In the Fig. 8 below calculating the dynamics of the NAD structure with time includes the angle between (point A~(PA,O3,PN), point B~N_{begin}, point C~N_{end}).




Fig. 8. The structure of NAD with highlighted atoms and a torsion angle.

Thus, the orientation features of the ADH enzyme's titratable amino acid residues have seen strongly correlating with the NAD co-enzyme's transformation. Thereby, the complete ADH+NAD

Disclaimer: This is not the final version of the article. Changes may occur when the manuscript is published in its final format.

structural behavior has to occur on the graphite surface during the adsorption processes. We have chosen the ambient temperature and water environment during the implementation of the MD numerical experiment; so far, the results in terms of pH and temperature coincide with data reported in the literature for the ADH enzyme in solution. Perhaps, above main results, Figs. 3b and 4b, representing the main discoveries or observations correlate well with experiments on the kinetics of the ADH enzyme. As can be seen from the MD results, presented in Figs. 3b and 4b, during the ADH+NAD adsorption the diffusion processes occur for 55 ns; after this period, ADH+NAD touch the graphite surface for a final relaxation on the surface. Following the adsorption of the ADH enzyme on the carbon surface, the coenzyme NAD from 55-60 ns begins to fluctuate, which is in full correlation with the opening of two catalytic loops associated with the NAD region. It is worth noting that current observation correlates with some experimental work related to the immobilization and adsorption of ADH enzyme on the carbon surface, the location of the enzyme and its fixation on carbon platforms. Moreover, the results also show that, in order to maintain an environment in which the ADH enzyme display good activity and to provide conditions for future technological applications, physiological conditions and ambient temperature can satisfactorily be applied to an enzymatic system involving the dehydrogenase enzyme. At the same time, the choice of carbon surfaces and platforms is motivated by a good control of the location of enzymes on the surface with very low enzyme consumption. The kinetic rates obtained for the ADH enzyme fixed to the carbon surface indicates a decrease in activity after the immobilization and fixation process, a significant loss of enzymatic activity observed after immobilization, even though the affinity between the enzymes and their substrates and coenzymes persists. To date, the MD observations of the structural behavior of ADH+NAD, diffusion and adsorption processes correlate well with the kinetic rates obtained for an enzyme anchored by ADH on a carbon platform with various kinetic parameters (mainly the concentration of substrates and coenzyme), indicating the substantial changes (the loss or decrease) in the enzymatic activity of fixed enzymes after the immobilization process on the carbon surface. Judging by experiments and kinetic studies, this is due to a significant loss of enzymatic activity of ADH in the presence of a carbon surface, and the anchored enzyme should lose enzymatic activity. Thus, for the ADH enzyme and NAD coenzyme, the carbon platform has probably cause not only diffusion activity, but also some restriction in diffusion process, thereby making them difficult for the reduced form of the coenzyme to escape from the bulk solution.

Disclaimer: This is not the final version of the article. Changes may occur when the manuscript is published in its final format.

CONCLUSION

In conclusion, the structural conformations of the alcohol dehydrogenase (ADH) enzyme with its cofactor nicotinamide adenine dinucleotide (NAD) were studied on a graphite-like surface (C-surface) using the MD (molecular dynamics) modeling method. The enzyme ADH+NAD was modeled in an aqueous medium. The MD simulations, conducted using the AMBER-18 package (a fast implementation of the *pmemd.cuda* module), provide valuable statistical data on key structural aspects of ADH, particularly its conformational changes, including the rotation of the catalytic domain, the coenzyme binding domain, and rearrangements of the active center. This analysis is essential for understanding the atomic and molecular details of a catalytically active enzyme. The 100-ns dynamic simulations allowed us to track the conformational and rotational changes of ADH+NAD in an aqueous environment.

We observed a non-trivial phenomenon: the conformational changes in ADH were correlated with structural transformations in its cofactor, NAD. However, it remains unclear whether the catalytic activity of ADH is preserved after such extensive conformational changes. Furthermore, our findings align with experimental studies on enzyme immobilization and adsorption onto carbon surfaces, as well as enzyme fixation on carbon platforms and oxide surfaces. The MD results of the distance and angle with time, appear to correlate well with experimental data on ADH+NAD kinetics where the ADH+NAD diffusion process occurs over approximately 50–60 ns, after which the complex reaches the graphite surface and undergoes final relaxation. Following ADH adsorption onto the carbon surface, NAD begins to fluctuate between 55 and 60 ns, aligning with the opening of two catalytic loops associated with the NAD-binding region. Experimental and kinetic studies suggest that this structural change significantly impacts enzymatic activity, leading to the loss of catalytic function in the immobilized enzyme. For ADH and NAD species on carbon platforms, diffusion restrictions may arise, potentially preventing the reduced form of the coenzyme from flowing freely in the bulk solution. The ADH+NAD complex was preserved in the simulations, with the cofactor modeled within the enzyme structure as in the original PDB file. Interestingly, NAD can be efficiently stabilized outside the enzyme, maintaining its catalytic efficiency in a selected NAD-enriched solution, a specially designed buffer or solvent system to stabilize and preserve its activity.

List of Abbreviations

- MDs: Molecular dynamics simulations
- ADH: Alcohol dehydrogenase
- NAD: Nicotinamide adenine dinucleotide

Disclaimer: This is not the final version of the article. Changes may occur when the manuscript is published in its final format.

- VMD: Visual Molecular Dynamics
- HIS: Histidine
- ASP: Aspartate
- GLU: Glutamate

- **Authors' Contributions,:**

- Conceptualization and supervision, P.P.G. and Kh.T.Kh.; methodology, Kh.T.Kh., M.A.I. and H.G.E.; validation, Kh.T.Kh., P.P.G., I.A.B.; formal analysis, Kh.T.Kh, M.A.H., I.A.B.; investigation, Kh.T.Kh., I.A.B., M.A.H.; resources, Kh.T.Kh., I.A.B.; data curation Kh.T.Kh, I.A.B., P.P.G.; writing—original draft preparation, Kh.T.Kh., I.A.B., P.P.G.; writing—review and editing, Kh.T.Kh., P.P.G., I.A.B., H.G.E., M.A.I.; project administration, P.P.G. and Kh.T.Kh.; funding acquisition, P.P.G. and H.G.E. All authors have read and agreed to the published version of the manuscript.

Availability of Data and Materials

Data supporting the results of this study are available upon request from the corresponding author.

Consent for Publication:

Not applicable.

Conflict of Interest

The author(s) declare no conflicts of interest regarding this manuscript.

Acknowledgments

The work has been performed within the framework of the joint research program of the JINR - Dubna State University. The work was performed within the framework of the state assignment of the Ministry of Science and Higher Education of the Russian Federation (No. 1024011000011-7-1.4.2;3.5.2 Conjugates boron-containing quantum dots with biovectors for the diagnosis and boron-neutron capture therapy of superficial malignant tumors (FEEM-2024-0011)). The MD calculations were performed on the servers of the Heterogeneous HybriLIT platform of the Multifunctional Information and Computing Complex (IVC), MLIT (Laboratory of Information Technology named after M.G. Meshcheryakov), JINR (Joint Institute for Nuclear Research).

Funding: This work was supported by funding from the JINR - ASRT (ARE) grant "Analysis of molecular modeling of the effect of nanometallic oxides on biological molecules". The work was carried out with the support of the Ministry of Science and Education of the Republic of Tajikistan

Disclaimer: This is not the final version of the article. Changes may occur when the manuscript is published in its final format.

within the framework of the project: "Molecular orientation of DNA on biocompatible metal-oxide materials" (Grant No.0122TJ1327).

REFERENCES AND NOTES

1. Wagenknecht PS, Penney JM and Hembre RT, Transition metal-catalyzed regeneration of nicotinamide coenzymes. *Organometallics* 22: 1180–1182 (2003).
2. Nakamura K and Yamanaka R, Light mediated cofactor recycling system in biocatalytic asymmetric reduction of ketenes. *Chem Commun* 16: 1782–1783 (2002).
3. Dmitry S. Bilan, Vsevolod V. Belousov. Genetically encoded probes for NAD⁺/NADH monitoring in Shemyakin-Ovchinnikov Institute of Bioorganic Chemistry, Moscow, Russia *Free Radical Biology and Medicine*. Volume 100, November 2016, Pages 32-42
<http://dx.doi.org/10.1016/j.freeradbiomed.2016.06.018>
4. A. Vrsalović Presečki, Đ. Vasić-Rački. Modelling of the alcohol dehydrogenase production in baker's yeast. *Process Biochemistry*. Volume 40, Issue 8, July 2005, Pages 2781-2791.
5. Tušek A, Šalić A, Kurtanjek Ž and Zelić B, Modelling and kinetic parameter estimation of alcohol dehydrogenase catalyzed hexanol oxidation in a microreactor. *Eng Life Sci* 12: 49–56 (2012).
6. Orlich B, Berger H, Lade M and Schomäcker R, Stability and activity of alcohol dehydrogenases in W/O-microemulsions: enantioselective reduction including cofactor regeneration. *Biotechnol Bioeng* 70: 638–646 (2001).
7. Sarah Höhn, Kai Zheng, Stefan Romeis, Martin Brehl, Wolfgang Peukert, Dominique de Ligny, Sannakaisa Virtanen, Aldo R. Boccaccini. Effects of Medium pH and Preconditioning Treatment on Protein Adsorption on 45S5 Bioactive Glass Surfaces // *Adv. Mater. Interfaces* 2020, 7, 2000420. doi.org/10.1002/admi.202000420
8. Tomás E. Benavidez, Daniel Torrente, Marcelo Marucho, Carlos D. Garcia. Adsorption and catalytic activity of glucose oxidase accumulated on OTCE upon the application of external potential // *Journal of Colloid and Interface Science*, Volume 435, 2014, Pages 164-170, ISSN 0021-9797, <https://doi.org/10.1016/j.jcis.2014.08.012>.
9. Fei Wang, Yu-Qing Zhang. Protein and Peptide Nanoparticles for Drug Delivery // *Advances in Protein Chemistry and Structural Biology*, 2015.
10. Welborn V. V. Structural dynamics and computational design of synthetic enzymes // *Chem Catalysis*. – 2021.
11. Norde W., Lyklema J. Why proteins prefer interfaces // *Journal of Biomaterials Science, Polymer Edition*. – 1991. – T. 2. – №. 3. – C. 183-202.

Disclaimer: This is not the final version of the article. Changes may occur when the manuscript is published in its final format.

12. Andrade J. D. (ed.). Surface and Interfacial Aspects of Biomedical Polymers: Volume 1 Surface Chemistry and Physics. – Springer Science & Business Media, 2012.
13. Gladyshev P. P., Shapovalov Yu. A., Kvasova V. P.. Reconstructed oxidoreductase systems. – NAUKA (The science). KazSSR, 1987.
14. Gladyshev P. P., Goryaev M. I., Shpilberg I. G., Shapovalov Yu. A. Sorption immobilization of NAD-dependent enzyme systems. I. The influence of electrostatic interactions on the orientation of alcohol dehydrogenase on the surface of the carrier. *Molecular Biology*, 1982, v. 16, no. 5, pp. 938-942 (in Russian).
15. Gladyshev P. P., Goryaev M. I., Shpilberg I. G. Sorption immobilization of NAD-dependent enzyme systems. II. The influence of hydrophobic interactions on the orientation of alcohol dehydrogenase on the surface of the carrier *Molecular Biology*, 1982, v. 16, no. 5, pp. 943-947. (in Russian).
16. Foresman J., Frish E. *Exploring chemistry // Gaussian Inc., Pittsburg, USA. – 1996. – T. 21.*
17. Leach A. R. *Molecular modelling: principles and applications. – Pearson education, 2001.*
18. D.A. Case, T.E. Cheatham, III, T. Darden, H. Gohlke, R. Luo, K.M. Merz, Jr., A. Onufriev, C. Simmerling, B. Wang and R. Woods. The Amber biomolecular simulation programs. *J. Computat. Chem.* 26, 1668-1688 (2005).
19. D.A. Case, H.M. Aktulga, K. Belfon, D.S. Cerutti, G.A. Cisneros, V.W.D. Cruzeiro, N. Forouzes, T.J. Giese, A.W. Götz, H. Gohlke, S. Izadi, K. Kasavajhala, M.C. Kaymak, E. King, T. Kurtzman, T.-S. Lee, P. Li, J. Liu, T. Luchko, R. Luo, M. Manathunga, M.R. Machado, H.M. Nguyen, K.A. O'Hearn, A.V. Onufriev, F. Pan, S. Pantano, R. Qi, A. Rahnamoun, A. Risheh, S. Schott-Verdugo, A. Shajan, J. Swails, J. Wang, H. Wei, X. Wu, Y. Wu, S. Zhang, S. Zhao, Q. Zhu, T.E. Cheatham III, D.R. Roe, A. Roitberg, C. Simmerling, D.M. York, M.C. Nagan*, and K.M. Merz Jr.* AmberTools. *J. Chem. Inf. Model.* 63, 6183-6191 (2023).
20. Tai-Sung Lee, Dan Mermelstein, Charles Lin, Scott LeGrand, Timothy J. Giese, Adrian Roitberg, David A. Case, Ross C. Walker* & Darrin M. York*, "GPU-accelerated molecular dynamics and free energy methods in Amber18: performance enhancements and new features", *J. Chem. Inf. Mod.*, 58:2043-2050 2018. 10.1021/acs.jcim.8b00462
21. Vinícius Wilian D. Cruzeiro, Marcos S. Amaral, and Adrian E. Roitberg, "Redox Potential Replica Exchange Molecular Dynamics at Constant pH in AMBER: Implementation and Validation", *J. Chem. Phys.*, 149, 072338 (2018). DOI: <https://doi.org/10.1063/1.5027379>.

Disclaimer: This is not the final version of the article. Changes may occur when the manuscript is published in its final format.

22. Kholmirdzo Kholmurodov (Ed.), "Models in Bioscience and Materials Research: Molecular Dynamics and Related Techniques", Nova Science Publishers Ltd., 2013, 219 p., ISBN: 978-1-62808-052-0.
23. Kholmirdzo Kholmurodov (Ed.), "Computational Materials and Biological Sciences", Nova Science Publishers Ltd., 2015, 208 p., ISBN: 978-1-63482-541-2.
24. DOI: <https://doi.org/10.2210/pdb3COS/pdb> (PDB ID: 3COS)).
25. Kholmurodov Kh.T., Baigunov I.A., Gladyshev P.P., Elhaes H. and Ibrahim M. "The Molecular Dynamics and Experimental Studies of the Structural Behavior of Alcoholdehydrogenase Enzyme on the Graphitic Sorbent Surfaces". *Res Dev Material Sci.* 20(2). RDMS. 000982. 2024. Published in Crimson Publishers (Volume 20 - Issue 2, June 2024).
26. Neto S.A., Forti J.C., Zucolotto V., Ciancaglini P., De Andrade A.R. "The kinetic behavior of dehydrogenase enzymes in solution and immobilized onto nanostructured carbon platforms". *Process Biochemistry*, 46 (2011), 2347–2352. <https://doi.org/10.1016/j.procbio.2011.09.019>
27. J.C. Forti, S. Aquino Neto, V. Zucolotto, P. Ciancaglini, A.R. De Andrade. "Development of novel bioanodes for ethanol biofuel cell using PAMAM dendrimers as matrix for enzyme immobilization". *Biosens Bioelectron*, 26 (2011), pp. 2675-2679.
28. D. Sokic-Lazic, S.D. Minter. "Citric acid cycle biomimic on a carbon electrode". *Biosens Bioelectron*, 24 (2008), pp. 939-944.
29. Radović, M., Hok L., Panić M., Cvjetko Bubalo M., Vianello R., Vinković M. and Radojčić Redovnikovića I. "Deep eutectic solvents as a stabilising medium for NAD coenzyme: unravelling the mechanism behind coenzyme stabilisation effect". *Green Chem.*, 2022, 24, 7661-7674. <https://doi.org/10.1039/D2GC02656B>.

Supplementary Materials

Movies S1-S2

Top and side views of the long 100-ns adsorption process of the alcoholdehydrogenase (ADH) enzyme with its co-factor nicotinamide adenine dinucleotide (NAD) on a graphite-like surface (C-surface) simulated by the MD (molecular dynamics) simulation method.

https://drive.google.com/file/d/1X5HdPXt4f14UR0Z_INcGjFrud5je2-9o/view?usp=sharing
https://drive.google.com/file/d/1atmztNahAKuczSB_KYr14p9si46IT9Lh/view?usp=sharing