



A Review on the Cell-Penetrating Peptides

Neelabh Datta 

School of Biology, Indian Institute of Science Education and Research Thiruvananthapuram (IISER TVM), Thiruvananthapuram, Kerala, India;
neelabhdatta@gmail.com

ORCID ID: 0000-0002-1577-5461

Abstract

The therapeutic potential of biomacromolecules in treating various diseases is undeniable; however, their application is significantly limited by the inherent barrier function of the cell membrane. Cell-penetrating peptides (CPPs), a class of small peptides comprising fewer than 30 amino acids, have garnered considerable attention due to their exceptional ability to traverse cell membranes. These peptides facilitate the intracellular delivery of various macromolecules, such as proteins, polypeptides, and nucleic acids, thereby overcoming the limitations imposed by the plasma membrane. CPPs exhibit versatile translocation properties, allowing them not only to penetrate cells independently but also to serve as delivery vehicles for therapeutic agents or CPP/cargo complexes. This unique capability has positioned them as promising candidates in the treatment and diagnosis of diseases, including cancer. Their ability to transport biologically active molecules across the plasma membrane enhances the efficacy of therapeutic interventions that would otherwise face significant challenges due to poor cellular uptake. This review explores the classification of CPPs based on their structure and properties, shedding light on the mechanisms that enable their efficient membrane penetration. It also touches into various therapeutic applications of CPPs, emphasizing their potential to revolutionize drug delivery systems. Despite their promise, the clinical utility of CPPs is hindered by challenges such as stability, specificity, and potential cytotoxicity. Addressing these limitations is crucial to unlocking the full potential of CPPs as innovative delivery tools in modern medicine.

Keywords

Cell-penetrating peptides; structure; detection; application; cancer therapy;

Introduction

Polypeptides, proteins, and genetic biomacromolecules have demonstrated significant potential in the treatment of various diseases. Despite their efficacy, their large molecular size and low lipophilicity hinder their ability to traverse cell membranes efficiently. This limitation restricts the therapeutic applications of these biomacromolecules, particularly in the medical field. Several techniques, including liposome-assisted delivery, viral vectors, electroporation, and microinjection, have been employed to facilitate their transport into cells. However, these methods often suffer from drawbacks such as low delivery efficiency, cytotoxic effects, and compromised cellular health. Poor targeting specificity has further limited their large-scale clinical use. As a result, the development of safe and efficient transmembrane delivery systems for biomacromolecules remains a critical challenge. Among the emerging solutions, cell-penetrating peptides (CPPs) have shown promise in addressing these issues.

One of the earliest discoveries in this field was the transactivator of transcription (TAT) protein derived from the human immunodeficiency virus (HIV), which was identified by Green et al. in 1988 to possess the ability to cross cell membranes [1]. Subsequent studies, such as those conducted by Brigati et al. in 2003, established that the amino acid residues located at positions 47–57 of the TAT protein are crucial for its transmembrane activity [3]. CPPs, also known as protein transduction domains (PTDs) or membrane transduction peptides (MTPs), are relatively small peptides consisting of 5–30 amino acid residues. These peptides have a unique capacity to cross cellular membranes independently and can also facilitate the transport of biomacromolecules through covalent or non-covalent interactions. Particularly CPPs exhibit remarkable versatility, as they can penetrate a wide variety of cell types and, when administered in vivo, are even capable of crossing the blood-brain barrier (BBB) [2]. To date, numerous naturally occurring and synthetically engineered CPPs have been identified. Examples include the TAT protein from HIV, the ANTP protein derived from the

Drosophila transcription factor, VP22 from herpes simplex virus type 1 (HSV-1), and synthetic peptides such as polyarginine and polylysine [2]. These peptides have garnered significant interest due to their broad range of transmembrane targets and their potential applications in therapeutic delivery. This review delves into the classification, structure, and therapeutic potential of CPPs, highlighting their diverse roles in modern medicine. It also examines their shortcomings and the challenges associated with their clinical application. Despite their immense potential, CPPs face limitations such as stability concerns, off-target effects, and potential toxicity, which must be addressed to fully realize their therapeutic value.

Structural characteristics and classification of cell-penetrating peptides

CPPs encompass a wide variety of types and are categorized based on factors such as their origin, mechanisms of uptake, biological applications, and physical properties. Despite these classifications, there is no universal consensus regarding a standardized grouping system. Typically, CPPs are short peptides consisting of 10 to 30 amino acids, often enriched with positively charged residues like lysine and arginine. These cationic amino acids contribute to their overall positive charge, facilitating interactions with the negatively charged phospholipid bilayer of cell membranes. Prominent examples of CPPs include TAT, penetratin, and transportan. CPPs are often grouped into three categories based on their chemical and physical characteristics: cationic, amphiphilic, and hydrophobic. Mainly cationic and amphiphilic CPPs account for approximately 85% of the total identified CPPs, while hydrophobic peptides constitute only about 15% [4]. Among these, cationic CPPs have been the most extensively studied due to their efficient interaction with cellular membranes. Figure 1 below shows the different types of CPPs available.




Figure 1: Classification and Properties of Cell-Penetrating Peptides (CPPs). Adapted from Gori et al. (2023).

<https://doi.org/10.1002/cmdc.202300236>

1. Cationic CPPs

Cationic CPPs, such as TAT, penetratin, polyarginine,

P22N, DPV3, and DPV6, are composed of peptide sequences that are rich in arginine, lysine, and histidine. The guanidinium group present in arginine plays a crucial role in forming hydrogen bonds with the negatively charged phosphate groups on cell membranes. This interaction facilitates CPP entry into cells under physiological pH conditions. These peptides often interact electrostatically with glycoproteins on the cell surface and are internalized independently of receptor-mediated mechanisms. The efficiency of cationic CPPs is influenced by the number and spatial arrangement of positively charged arginine residues within their structure [5]. Most cationic CPPs contain more than five positively charged amino acids, which are critical for their functionality [6]. Among these, polyarginine sequences show the highest efficiency for cell entry, making them particularly promising for therapeutic applications [7]. Studies by Chu et al. (2015) demonstrated that the internalization potential of oligoarginine increases with peptide length, with R8 to R10 being the optimal range for delivery applications [8]. Experimental analyses of oligoarginines ranging from three to twelve residues revealed that efficient membrane perforation requires at least eight arginine residues, with activity progressively increasing as the number of residues increases [2]. Although lysine shares a positive charge with arginine, it lacks the guanidinium group, resulting in comparatively lower membrane permeability when used in isolation [9]. Research by Futaki et al. (2001) further underlined that cationic CPPs with at least eight positively charged residues achieve superior membrane-penetrating efficacy [10].

Other amino acids also play critical roles in CPP functionality. For example, substituting tryptophan at position 14 with phenylalanine in penetratin, results in a loss of membrane permeability [11]. Nuclear localization signal (NLS) sequences represent a subset of cationic CPPs. These are short peptides enriched in arginine, lysine, and proline that facilitate transport to the nuclear pore complex. NLS peptides are further classified as mono- or bi-partite based on whether they contain one or two clusters of basic amino acids. An example of a monotype NLS is PKKKRKV from Simian virus 40 (SV40), while bitype sequences are exemplified by nucleoplasmin. The smallest known NLS sequence with membrane-permeating ability is KRPAAT-KKAGQAKKKL [2]. However, due to their lower charge (fewer than eight positive residues), NLS peptides are not inherently effective CPPs [12]. When conjugated with hydrophobic sequences, NLS peptides can form amphiphilic CPPs with enhanced properties. Cationic CPPs are being actively explored for their versatility in biomedical applications, including drug delivery, gene therapy, and molecular imaging. These peptides represent a promising avenue for advancing therapeutic strategies by enabling efficient intracellular transport of therapeutic molecules.

2. Amphiphilic CPPs

Amphiphilic cell-penetrating peptides (CPPs) are a diverse group characterized by their ability to possess both hydro-

philic and hydrophobic properties. They are further subdivided into four categories: primary amphipathic, secondary α -helical amphipathic, β -sheet amphipathic, and proline-rich amphipathic CPPs [13]. Primary amphipathic CPPs are either derived from natural protein sequences or designed by combining hydrophobic peptide domains with nuclear localization signals (NLSs). For instance, MPG (GLAFLGFLGAAGSTMGAWSQPKKKRKV) and Pep-1 (KETWWETWWTEWSQPKKKRKV) are synthetic examples of this. Both are based on the NLS sequence PKKKRKV from Simian Virus 40 (SV40) but incorporate hydrophobic domains to enhance their amphipathic nature. In MPG, the hydrophobic domain is derived from a fusion sequence, while in Pep-1, it is represented by the peptide segment KETWWETWWTEW. These hydrophobic sequences are joined to the NLS through a short linker peptide, WSQP [14]. Alternatively, some primary amphipathic CPPs originate from naturally occurring proteins. Examples include pVEC, ARF(1-22), and BPrPr (1-28), which demonstrate similar structural and functional features but are directly sourced from protein sequences rather than synthetic design. Secondary α -helical amphipathic CPPs function by interacting with the cell membrane through their α -helical conformation. These peptides possess distinct hydrophilic and hydrophobic regions distributed across different faces of the helix, allowing effective membrane association and penetration. A representative example is MAP (KLALKLALKALKAAALKLA), which demonstrates significant transmembrane activity through this mechanism [2]. The functionality of β -sheet amphipathic CPPs is closely tied to their ability to form β -sheet structures. The alignment of amino acid residues in pleated β -sheets is essential for their interaction with and entry into cell membranes. Peptides unable to establish this configuration lose their membrane-permeating capabilities, highlighting the critical role of β -sheet formation in their mechanism of action [15].

Proline-rich amphipathic CPPs are characterized by their high proline content, which facilitates the adoption of a polyproline II (PPII) helical structure in aqueous environments. Unlike the typical right-handed α -helix with 3.6 residues per turn, PPII helices are left-handed and consist of approximately three residues per turn. Examples of proline-rich amphipathic CPPs include the bovine antimicrobial peptide Bac7 and synthetic peptides such as (PPR) $_n$, where "n" can vary between 3 and 6 [2]. These unique structural features contribute to their efficient cellular uptake and potential therapeutic applications.

3. Hydrophobic CPPs

Hydrophobic CPPs are composed predominantly of nonpolar amino acid residues and possess a net positive charge of less than 20%. Their functionality relies on the presence of hydrophobic motifs or specific chemical groups essential for traversing the lipid bilayer. Despite being less extensively studied than other CPP classes, hydrophobic CPPs have notable examples, such as fibroblast growth

factor-derived peptides like K-FGF and F-GF12, which are associated with Kaposi's Sarcoma [16]. These peptides represent an alternative strategy for intracellular delivery, particularly in hydrophobic environments where traditional CPPs might face limitations.

Membrane penetration mechanism of cell-penetrating peptides

Despite the fact that CPPs have emerged as a research hotspot in recent years, their membrane-penetrating mechanism is still contemplated. On the internalisation methods of several CPPs or CPP/cargoes, there appears to be unanimity [17,18]. At the moment, there are primarily two mechanisms of CPPs transmembrane: direct transmembrane (energy-independent pathway) and energy-dependent endocytosis depending on whether or not energy is needed during internalisation [19]. The specific mode of entry into cells is linked to the physical and chemical properties of CPPs, the size, charge, and type of cargo transported, the accumulation of CPPs and cargo complexes, the medicated tissues and cell types, and so on [20] (Figure 2).




Figure 2: Mechanisms of Cellular Uptake of CPPs. This figure depicts the two primary pathways by which CPPs enter cells: endocytosis and direct translocation. Endocytic routes include macropinocytosis, clathrin-dependent, caveolae-mediated, and clathrin/caveolin-independent mechanisms, often requiring vesicular trafficking. Direct translocation involves pore formation (barrel stave, toroidal) or membrane disruption via inverted micelle and carpet models. Figure adapted from Trabulo, S et al. (2010). <https://doi.org/10.3390/ph3040961>

1. Direct membrane penetration

Direct membrane penetration is an energy-free pathway, with proposed mechanisms including inverted micelle mode [21], carpet mode [22], perforation mode i.e. "Barrel-Stave" model [23], and plasma membrane sparse mode [20]. Each one of these modes necessitate cationic CPPs to

first engage with anionic components on the plasma membrane, such as phospholipid bilayers, causing modifications in plasma membrane integrity, and the successive internalisation process is dependent on the type and concentration of CPPs, the quantity of cargo, etc. attributes, cell lines treated, and storage conditions, etc [2]. While these mechanisms can describe some facets of CPP transmembrane transport, none can establish a comprehensive internalization pathway relevant to all types of CPPs. When using nuclear magnetic resonance (NMR) to study the interaction between penetration and phospholipid membranes, the inverted micelle model was proposed [24]. The cationic CPPs first interact with the anions on the phospholipid membrane, and then the phospholipid bilayer rearranges and shuttles due to the interaction between the CPPs' hydrophobic amino acid residues and the phospholipid membrane, and finally the CPPs are wrapped [25]. The micelle, entombed within the inverted micelle, moves to the other side of the lipid bilayer membrane and explicitly discharges CPP into the cytoplasm [2]. The mechanism cannot work for CPPs without hydrophobic amino acid residues such as TAT and oligoarginine, since this model assumes hydrophobic amino acid residues are required to form flipped micelles. Carpet structures have been employed to characterize not only the translocation mechanism of some antimicrobial peptides, but also the cytotoxicity of CPPs at high concentrations. Throughout this mode, cationic CPPs carpet the surface of the negatively charged plasma membrane like on flooring, and then the hydrophobic amino acid residues engage with the plasma membrane's hydrophobic core because the basic amino acid residues in the CPPs face the plasma membrane surface. When CPP concentrations reach a certain level, the phospholipid membrane becomes regionally compromised, permitting CPPs to penetrate the cell [26, 27]. The punch sequence, also known as the barrel stave sequence, is a mechanism for antimicrobial peptide internalisation in bacteria. The α -helical structure of the amphiphilic CPPs is required for the formation of the pore-type channel. CPPs are grouped together on the cell surface and implanted in parallel structures into the cell membrane to establish a barrel-like conduit [17]. The hydrophobic area of the α -helix adheres to the membrane lipid's outer surface to construct the conduit, while the hydrophilic region binds to the phospholipid's hydrophilic head, forming the conduit's central core and facilitating CPP transmembrane translocation and internalisation. The CPPs are delivered into the cytoplasm [18]. The "plasma membrane thinning" impact was first used to explain magainin's mechanism of action [28]. After CPPs form a carpet structure, the anomaly of the interaction between the outer leaflet charges induces the rearrangement of negatively charged membrane lipids and the thinning of the plasma membrane in this model. CPP interaction or aggregation on the plasma membrane surface lowers the surface tension of the plasma membrane's local surface, enabling CPPs to insert into the plasma membrane and penetrate the cell [2]. However, since no plasma membrane permeability

was observed, this mechanism is highly unlikely to be plausible. It is possible to explain why the cell membrane does not appear permeable by considering the transient properties of plasma membrane pores combined with the repair response of the cell membrane [29].

2. Endocytic pathway

The involvement of endocytic pathways in cell-penetrating peptide (CPP) internalization has been a central focus in understanding the mechanisms of CPP-mediated membrane penetration. Initial studies suggested that CPP uptake occurred through a non-receptor-mediated, non-transporter-dependent, and energy-independent direct internalization process. Over time, however, four distinct endocytic pathways have been identified: clathrin-mediated endocytosis [30], caveolin-mediated endocytosis [31], macropinocytosis [32], and clathrin- and caveolin-independent endocytosis [33]. Early research by Vives et al. (1997) demonstrated that TAT could enter cells at 4°C or under energy-depleted conditions, supporting the idea of energy-independent internalization and excluding the involvement of endocytosis [34]. However, subsequent studies revealed limitations in these early methods, such as the use of methanol or formaldehyde for cell fixation, which could redistribute CPPs on the cell surface, thereby obscuring the true internalization process [35]. Cationic CPPs also exhibit strong membrane affinity, complicating the distinction between membrane-bound and internalized peptides when analyzed by flow cytometry. This limitation often led to overestimation of internalization rates [2]. To address these concerns, Richard et al. (2003) utilized more precise techniques, confirming through flow cytometry and cell fixation that TAT internalization occurs via endocytosis [36]. Similarly, Wadia et al. (2004) observed colocalization of TAT-Cre with the endocytic marker FM4-64, providing direct evidence for an endocytic mechanism underlying TAT-mediated transport of exogenous molecules [32].

While some CPPs can bypass membranes through direct translocation, the majority rely on endocytic processes for internalization. This mode of entry, however, poses challenges for cargo release, as CPP-bound materials often remain sequestered in endosomes and are subsequently degraded in lysosomes unless they successfully escape the endosomal compartment [37, 38]. In receptor-mediated endocytosis, CPPs or their cargo bind to membrane receptors, inducing curvature through interaction with epsin proteins. Clathrin and hetero-tetrameric adaptor proteins (AP-2) assemble to form clathrin-coated pits, which mature into vesicles encapsulating the CPP-cargo complex [33, 39]. The Caveolin-mediated endocytosis pathway involves CPP binding to receptors located in lipid rafts—hydrophobic membrane regions enriched in sphingomyelin and cholesterol. The cavin-1 and caveolin proteins facilitate vesicle formation and endosome generation, mediating

cargo internalization [40, 41]. The preference for caveolin-mediated pathways may vary based on the size and properties of the associated cargo [42].

When CPPs are bound to larger macromolecular cargos (e.g., >30 kDa), they often employ macropinocytosis, an actin-dependent endocytic pathway characterized by membrane ruffling and vesicle formation. This process is initiated by CPP interactions with membrane proteoglycans, which activate rac proteins. These proteins, in turn, trigger F-actin rearrangements that facilitate vesicle formation and cargo uptake [33, 43]. Unlike other pathways, macropinocytosis is receptor-independent and often lipid raft-dependent, making it particularly suited for CPPs carrying large payloads [2, 44]. In specialized cells like macrophages, CPP uptake can occur through an alternative mechanism that does not involve clathrin or caveolin. Instead, CPPs are opsonized and recognized by cell surface receptors such as Fc receptors. Actin activation drives the internalization of CPP-cargo complexes into membrane-coated vesicles [45]. Regardless of the endocytic pathway utilized, CPPs and their cargos must escape endosomes to avoid lysosomal degradation and exert their biological effects. This escape is facilitated by processes such as pH gradient alterations, vesicle accumulation, and CPP interaction with charged endosomal membranes. These interactions can lead to increased membrane rigidity and eventual rupture, enabling the release of CPPs and their associated cargos into the cytoplasm [5, 6]. The choice of endocytic pathway for CPP internalization is influenced by the peptide's physicochemical properties and the characteristics of the bound cargo, including size, charge, and hydrophobicity. For example, TAT has been shown to use different pathways—lipid raft-mediated endocytosis for protein cargos and clathrin-mediated endocytosis for smaller fluorescent molecules. Understanding these mechanisms is crucial for optimizing CPP-based delivery systems for therapeutic applications.

Factors affecting the cellular uptake mechanism

Despite the fact that various CPPs share a lot of commonalities, their absorption methods may differ greatly. This results in inconsistent findings, mostly due to the numerous variables that influence the cellular absorption and translocation process. The physicochemical characteristics, concentration of the peptide and its payload, and features of the plasma membrane, including its lipid and protein composition, may be broadly categorised as variables that affect the absorption pathways of cell penetrating peptides. Because to variations in experimental settings, conflicting data about the internalisation process of CPPs frequently occur. The CPP concentration is the primary critical element. It has been demonstrated in several instances that the administered concentration significantly influences the uptake mechanism. The net charge of the peptides is an-

other consideration, particularly the positive charges that result from arginine residues [46]. The majority of CPPs include arginine residues, which are better for the transport and absorption of CPP than lysine (in particular, the guanidinium group of arginine) [46]. Amphipathicity is another element that has been linked to absorption. Whereas non-amphipathic CPPs need endocytosis, primary and secondary amphipathic peptides can directly cross the cell membrane at low concentrations [47]. The internalisation process may also be affected by the experiment's temperature. The temperature dependency of R8 translocation through plasma membranes has been seen by Fretz et al., 2007 [48]. At 4 °C, diffuse signals from the fluorescently tagged peptide are more pronounced in the cytoplasm, which is often an indication of a direct translocation across the membrane, they have discovered [48]. At 37 °C, however, both diffuse and punctate signals were seen, suggesting that an endocytic process may be activated at higher temperatures [48]. The absorption route can also be significantly influenced by the cargo molecules coupled to CPPs.

1. Role of cargo molecules in CCP uptake

The cargo coupled to the CPP is frequently a crucial internalisation component. The absorption of TAT conjugated to peptides and globular proteins in live cells was compared by Tünnemann et al., 2006 [49]. They discovered that the uptake process was significantly influenced by the size of the payload fused to TAT. While the TAT-peptide conjugates were dispersed widely throughout the cell, the bigger complexes comprising proteins were visible inside vesicular structures [49]. This suggests that a separate absorption method results from the size of the cargo, which naturally determines the size of the total complex. The likelihood of the complex being taken up by direct translocation increases with decreasing size. Endocytosis, however, predominates at bigger dimensions. This is comparable to the situation where the presence of a cargo molecule chooses between two alternative endocytosis methods when comparing the uptake mechanism of unconjugated TAT versus TAT fused to a cargo [39]. Whereas its conjugated counterpart is more likely to adopt CvME, unconjugated TAT favours CME for cell entry [46]. Cargo impact can also affect how other arginine-rich peptides are taken up. The effects of cargo molecules on the absorption of R7 and R7W were examined by Maiolo et al., 2005 [50]. In the cytoplasm of the tested cells, the peptides alone displayed diffuse signals. The diffuse signal from endocytic vesicles was significantly reduced following fusing to cargo peptides, but the punctate signal barely changed.

2. Role of concentration in CCP uptake

Since that multiple uptake routes can be stimulated by CPP concentration, it is a crucial component. It is thought that endocytosis often takes place at low peptide concentrations and that direct penetration takes over at greater concentra-

tions. Fretz et al., 2007 [48] evaluated the dependency on concentration while examining the impact of temperature on the absorption of R8. Vesicular staining in the cytoplasm was seen at lower peptide concentrations, indicating endocytic absorption [46]. Vesicular and diffuse labelling were seen at greater concentrations, suggesting that direct penetration and endocytosis may take place concurrently [51]. The concentration-dependent absorption of R9 and TAT has also been investigated [51]. Moreover, these peptides exhibit substantial cytosolic labelling at higher concentrations and mostly vesicular signals at lower concentrations. Nevertheless, things become more difficult since endocytic inhibitors had only a minimal impact on the absorption of R9 and TAT at low concentrations (<5 uM), but clathrin inhibitors appear to have a significant impact at higher concentrations [46]. In addition to the quick, nonendocytic absorption via nucleation zones, it was shown that the peptides are, to some extent, taken up by vesicular structures at greater concentrations [52]. According to these findings, the impact of CPP concentration on the uptake process may be far more nuanced than first believed.

3. Role of the type of cells involved in CCP uptake

The CPP uptake process is greatly influenced by the cell lines that were employed in the research. The nature of the extracellular matrix and the plasma membrane in particular can have a significant impact on CPP uptake. It is well known that the extracellular matrix's negatively charged GAGs and the positively charged CPPs have their first interaction during internalisation. Hällbrink et al, 2004 looked at the impact of the peptide-to-cell ratio on the absorption of cells into Chinese hamster ovary (CHO) cells [53]. To be more precise, they were interested in how the absorption would alter if the number of peptides, rather than the concentration, were raised while maintaining the same number of cells. They also looked at how cell number and confluence affected uptake. They demonstrated that increasing the internal peptide concentration more effectively than increasing the exterior peptide concentration required double the incubation volume at a certain number of cells. Moreover, applying a constant peptide concentration to various cell densities showed that the uptake decreased with increased confluence. The variable membrane composition or the various endocytosis patterns of the developing cells may be the reason of this. Another hypothesis is that when confluence rises, membrane access diminishes [46]. These findings offer an illustration of how experimental variables in cell culture might affect uptake effectiveness. A thorough analysis of the absorption of 22 distinct CPPs in four different cell lines was conducted by Mueller et al., 2008 [54]. They investigated the internalisation of several of the most well-known CPPs, including penetratin, TAT, transportan, Pep-1, MPG, MAP, R7, and R9. They employed Cos-7, HEK293, HeLa, and MDCK as

representative cell lines. According to their behaviour, the data allowed for the classification of CPPs into three groups: high (penetratin, transportan, MAP), medium (TAT, Pep-1, MPG), and low cellular uptake [46]. The results demonstrate that some of the peptides are taken up by cells in a cell-dependent manner, which is interesting. For instance, Cos-7 cells, which resemble fibroblasts and were isolated from monkey kidney tissue, took up MPG preferentially [46]. This could be because the virus that was utilised to create the Cos-7 cell line, SV-40 big T antigen, includes a NLS that was generated from MPG [55].

A recent research by Gronewold et al., 2017 examined how cell lines affected the absorption of a CPP with potential anticancer action [56]. The antimicrobial peptide CAP18's C-terminal domain was used to create sC18, the CPP. In addition to HEK293, a noncancer cell line, its absorption was examined in the cancer cell lines HeLa, PC-3, HCT-15, and MCF-7 [56]. It's interesting to note that nuclear accumulation and a diffuse fluorescent signal with punctate distribution were seen in the cytoplasm of all cancer cell types. Nevertheless, only a punctate distribution and little to no peptide in the nucleus were found in the noncancer cell line.

Detection of CPPs by Fluorescence

Fluorescence is a widely utilised method for locating and investigating CPPs. CPPs should be conjugated to the appropriate fluorophore molecules in order to use fluorescence procedures on them. The fluorophore moiety often has a covalent link attached to CPP. In all fluorescence techniques, CPP linked to a fluorophore is first incubated with cells, vesicles, or other target structures to enable its penetration into the cell. The fluorophore in the target structure is ignited by the absorption of light after an appropriate amount of time has passed. The equipment detects the light that the fluorophore produces when it transitions back to its ground state. A certain frequency of light is emitted by each fluorophore. Depending on the molar extinction coefficient and quantum yield, the fluorophore's efficiency, the excitation light's intensity, and the fluorophore's efficiency determine how much light is emitted. But, the fluorophore's neighbourhood can affect how much light it emits, and from this information, it is possible to infer how the fluorophore interacts with other molecules like proteins, peptides, or lipids. The validation of the effective delivery of cargo by identifying the right physiological impact of the cargo in the cell is the best technique to identify the internalisation of the CPP-cargo construct into the cells as well as into its target component (organelle) [57]. Another strategy is to co-localize the fluorescence light that the CPP-delivered payload emits with the light that certain target markers (molecules, organelles, cell types, tissues, or organs) emit or absorb. Moreover, CPP assesses possible cytotoxicity in relation to construct concentration to ensure safe distribution. The precursor of

interleukin 1 alpha that is armed with a nuclear localization sequence (NLS) and connected to green fluorescent protein (GFP) has been demonstrated to possess CPP abilities using fluorescence microscopy and flow cytometry [57]. It delivers GFP into the nuclei of Jurkat and HeLa cells. It was also demonstrated that, following intraperitoneal treatment, it is capable of delivering proteins into the cells of many mouse organs, including the spleen, liver, and intestine [58]. Even at 100 μ M concentration of the construct, no cytotoxicity was seen. It has been proved that the whole Antennapedia homeodomain protein may preferentially enter the cells by macropinocytosis using the same combination of fluorescent techniques and low temperature [59]. He et al., 2013 originally discovered a number of unique CPPs using a mixture of fluorescence techniques [60]. These CPPs were chosen utilising synthetic membranes and named spontaneous membrane-translocating peptides (SMPTs).

Tetramethylrodamine (TAMRA) and Alexa Fluor 546 dye were able to enter the cytoplasm of several cells via these peptides when endocytosis was blocked [57]. In contrast to SMTP unconjugated TAMRA, which was quickly eliminated when administered through injection to mice, SMTPs transported TAMRA to several organs where it could be identified even two hours after treatment. They hypothesised that SMTPs may convey a wide variety of additional polar chemicals into cells. Although while the internalisation process of the CPP prototype TAT, which was the first CPP found, is not fully understood, it is currently being thoroughly researched. The action of TAT as a CPP is usually associated with the activity of other CPPs, and it also acts as a model and reference CPP [57, 61]. Fluorescence spectroscopy, flow cytometry, and confocal imaging were used to investigate the cell-uptake of fluorescein isothiocyanate (FAM) tagged TAT [62]. It was discovered that the TAT-FAM construct may enter cells without using endocytosis through two entrance pathways with various energy requirements. A lot of positively charged amino acids make up CPPs like TAT, which are known for their propensity to interact with negatively charged biological structures [57]. To increase their availability and blood circulation time, fewer cationic CPPs are required. The usage of membrane interacting proteins as the source of the prospective membrane transferring peptides was used to solve this issue. According to Kim et al., 2015, many hydrophobic CPPs may be generated from the sequence of annexin [61]. These CPPs outperformed the CPP prototype TAT as cargo-delivery agents, as demonstrated by fluorescence flow cytometry and other techniques. These peptides exhibited negligible immunogenicity, extremely mild cytotoxicity, and high serum stability.

Detection of CPPs by Radioactivity

CPPs can be labelled with various markers for cell imaging in addition to fluorophores. Labelling radioactive ma-

terial is one strategy. Radioactive detection of CPPs has several advantages over other detection methods. It is highly sensitive, allowing for the detection of very small amounts of CPPs, and can be used to track the movement of CPPs within cells or tissues. It is also relatively easy to perform and does not require specialized equipment. Cells are typically initially treated with labelled CPP in a normal experiment [57]. To remove the labelled CPP that has been adsorbed on the membrane's exterior, cells are next rinsed with a solution that includes a suitably diluted acid [57]. The amount of internalised CPP is then estimated after the radioactivity counter determines the radioactivity. The kinetics of CPP internalisation may be observed if the incubation of the cells is stopped at various time points. TP labelled with 125I is one of the original examples of radioactively labelled CPP [63]. In order to observe the dynamics of the tagged TP internalisation, the amount of internalised TP inside Bowes cells could be calculated as a function of time [57]. The difficulty of this approach to distinguish between intact radioactive CPP and its radioactive by-products after the cleavage in the cell is a disadvantage. Using 125I to mark the four cationic CPPs TAT, octa- and nona-arginines, and MAP and monitor their uptake into CHO cells is a more contemporary example [64]. To examine the internalisation of the construct in HeLa and breast cancer cells, other proteins labelled with 125I that are involved in the metabolism of arginine can be fused to CPPs [65]. The results of the first of these two investigations showed that whereas TAT and poly-arginines were primarily kept in the cytosol, MAP was able to cross the nuclear membrane [57]. It should be noted that the presence of Tyr in the peptide structure is a requirement for labelling with 125I; if this is not the case, Tyr may be added to extend the peptide's sequence, but its potential impact on CPP behaviour should be investigated independently [57].

This was accomplished in the work carried out by Zaro et al., 2009 using poly-arginines [64]. In recent radioactivity experiments after CPP internalisation, 68Ga [66] and 64Cu were also utilised as nuclides [67]. A study that indicates other nuclides that might be utilised for the radioactive labelling of CPPs, including 99mTc, 111In, and 177Lu, was recently published by Gharibkandi et al., 2020 [68]. This analysis does not particularly address CPPs, but is generally focused on the radioactive labelling of peptides. There are some limitations to the use of radioactivity for CPP detection. It requires careful handling and disposal of radioactive materials, and there are concerns about the potential health risks associated with exposure to radiation. Besides, some CPPs may not be compatible with radioactive labeling due to their chemical properties.

Application of cell-penetrating peptides as transport vehicles

In addition to neurological disorders, asthma, local ische-

mia, diabetes and cancer, the CPPs- dependent drug delivery system has been used to treat a wide variety of diseases. As an effective delivery tool, CPPs have been used to introduce cytotoxic drugs into tumor cells to induce apoptosis in tumor cells.

1. CPPs facilitate transmembrane transport of small molecule drugs

Because of their small size and lipophilicity, small molecule anticancer drugs can dissipate into tumour cells in efficient manner. Multidrug resistance (MDR) develops when tumour cells are subjected to the same drug continuously. To address this issue, some researchers have attempted to combine these drugs with CPPs in order to facilitate the entry of these small molecule drugs into cells [69]. Dubikovskaya et al. formed a R8-taxol covalent compound by disulfide bonding octamer arginine R8 to the anticancer drug paclitaxel [69]. The results revealed that the R8-taxol covalent compound had a similar effect to paclitaxel alone in taxol-sensitive tumour models, whereas in taxol-resistant tumour models, R8-taxol covalent Drugs are more likely to trigger tumour cell apoptosis than paclitaxel alone [2]. There are numerous advantages to combining small molecule drugs and CPPs. It can increase drug water solubility and improve drug utilisation rate, in addition to overcoming multi-drug resistance. Lee et al., 2011 used chemical methods to combine doxorubicin, TAT, and polymerized chitosan backbones to generate chitosan/doxorubicin/TAT chimaeras that were compatible with TAT-free chimaeras in order to enhance the cytotoxicity and targeted transport of anticancer drugs [70]. This chimaera has more efficient cell internalisation than doxorubicin or chitosan/doxorubicin, and it can change the distribution of doxorubicin in the organism, enhance tumour localization, and thus considerably constrain tumour growth [70].

2. CPPs facilitate transmembrane transport of peptides and proteins

Numerous cancers can be brought on by mutations in tumour suppressor genes. [71, 72]. To reinstate the activity of these proteins and accomplish the goal of cancer treatments, some researchers attempt to insert full-length proteins or polypeptides pertaining to tumour suppressor genes into cancer cells. In order to treat mice with advanced peritoneal cancer metastasis, Snyder et al. created a TAT-p53 chimeric peptide and administered it intraperitoneally. The average survival time of the mice in the experimental group was more than 6 times that of the control group, which had a mean survival time of 10 days [73]. Hosotani et al., 2002 covalently joined a short peptide made up of 20 amino acid residues of the p16 protein to penetratin by a disulfide bond in order to reinstate the functioning of the p16 protein, and they then tested its effectiveness in an animal model of pancreatic cancers [74].

According to the findings (tumor in treatment group: 79 ± 17 mg; tumor in control group: 149 ± 12 mg), the complex could greatly slow the growth of cancers [74]. Apoptosis induction of tumour cells was attempted by certain researchers utilising various therapeutic techniques. As a protein with mitochondrial origin, Second Mitochondrial-Derived Activator of Caspase (SMAC) is crucial for the regulation of apoptosis [75]. SMAC can deactivate apoptosis-inhibiting proteins when it is released from mitochondria, hence increasing apoptosis. The 7 amino acid residues at the amino terminals of SMAC and TAT were combined to create a chimeric peptide, which was identified by Fulda et al. Research conducted in vitro revealed that while this chimeric peptide was ineffective at stimulating tumour cell apoptosis, it did make tumour cells more susceptible to apoptosis effectors such tumour necrosis factor-related ligand that induces apoptosis (TNF-related apoptosis inducing ligand, TRAIL) [76]. The conjunction of TRAIL and SMAC-TAT peptide can greatly slow the formation of tumours in a mouse model of human glioma xenografts, and when 0.6 or 2ug of TRAIL is employed, tumour cells can be completely eliminated [76]. The ribosome inactivating protein (RIP) generated from plants known as gelonin can efficiently limit protein translation when its half maximum inhibitory concentration (IC₅₀) is at the picomolar level [2]. Poor membrane permeability makes it difficult to treat tumours and frequently fails [77]. A chemical combination of gelonin and TAT or LMWP was used by Park et al. to overcome this obstacle [78]. In mouse colon cancer cells, the gelonin chimeric peptide was shown to have strong antitumor activity when it was injected. At a dose of 100 ug, the chimeric peptide can completely inhibit tumor growth [78].

3. CPPs facilitate transmembrane transport of genes

For the treatment of cancer, genes, antisense oligonucleotide strands, and small interfering RNA (siRNAs) may be useful tools [79]. A particularly alluring therapeutic approach, RNA interference technology is extremely selective to target genes and has a special tumour suppressor impact. Drugs containing siRNA have a minimal ability to penetrate cells due to their enormous size, strong charge, and size. CPPs have drawn a lot of attention due to their low toxicity in endeavors to facilitate siRNA entry into cells using cationic liposomes, cationic polymers, and CPPs. In a covalent or non-covalent manner, siRNA can attach to CPPs. Despite the fact that covalent bonding is the best type of binding, tight complexes and polymers are simple to produce because of the strong charge interaction between the two [2]. It is more difficult to cross the cell membrane than CPP when combined with another molecule in a 1:1 covalent manner [2]. Cationic CPPs' charge can be easily neutralised by negatively charged siRNA, rendering both siRNA and CPPs ineffective for their intended biological purposes. Researchers have decided to

combine siRNA with CPPs in a non-covalent form because to the several disadvantages of covalent binding. They are mostly focused on the viability of siRNA-CPP chimeric peptide research in vitro, however animal research is also ongoing. In the angiogenesis requisite for tumour growth, vascular endothelial growth factor (VEGF) plays a significant role.

The VEGF gene can be silenced or VEGF receptors blocked to treat malignancies. A cholesterol/R9/siVEGF combination was created by Kim et al., 2006 using R9 chimaera as a carrier and anti-VEGF siRNA (siVEGF) to drastically reduce tumour development and angiogenesis [80]. Cyclin B is a mitotic cycle protein that is necessary for cell division, but tumour cells manifest uncontrolled expression of this protein [2]. To reach therapeutic goals, several researchers try to use RNA interference technologies. Cromez et al. formed an anti-cyclinB1 siRNA complex by combining anti-cyclinB1 siRNA with the shortened version of MPG (MPG-8), which can significantly inhibit cell proliferation by inhibiting the expression levels of cyclinB1 and causing cell cycle arrest [81].

4. CPPs-facilitated transmembrane transport of nanoparticles

Cell-penetrating peptides have been extensively used in the transmembrane transport of numerous nanoparticles, including magnetic nanoparticles, lipid nanoparticles, gold nanoparticles, micelles, and quantum dots, despite the fact that the transmembrane mechanism of these molecules is not fully understood. TAT and cross-linked iron oxide particles (CLIOs) were coupled by Josephson et al., 1999 to create a TAT/CLIO complex with an average particle size of 41 nm [82]. The efficiency of the transmembrane was higher than CLIO without TAT. It is advantageous for MRI or magnetic sorting since it is approximately 100 times higher and can mark cells effectively [82]. The transmembrane efficiency of TAT/CLIO increased nonlinearly as the ratio of TAT/CLIO increased, and when the ratio reached 15, the highest transmembrane efficiency was found [2].

The complex created when liposomes and CPPs combine may efficiently enter a variety of cells, including mouse heart muscle cells, mouse lung cancer cells, and human breast cancer cells, and the effectiveness of liposomes' membrane penetration is related to the density of CPPs [83]. The penetratin/liposome complex has a higher rate of membrane penetration than the TAT/liposome complex and reaches its maximum absorption in under one hour. Researchers are very interested in employing CPP/nanoparticle complexes to transport diverse anti-cancer medicines since CPPs can successfully facilitate the transmembrane transport of different nanoparticles. Paclitaxel, a drug that can effectively enter malignant glioma cells and suppress tumour cell proliferation, was loaded into lipid nanocapsules with CPP modifications by

Balzeau et al., 2013 [84].

5. CPPs role in Cancer Therapy

One challenge in treating cancer was that tumour microenvironment or other obstacles prevented medication transport to tumour cells, particularly in cases of duodenal and brain gliomas. CPPs provided a fresh viewpoint on how to get through a semipermeable hydrophobic barrier and achieve excellent drug delivery in tissue and subcellular architecture. The majority of CPPs reacted with the high-density anionic charge on cell membranes and possessed positive side chains. Polyarginines of various lengths were often employed for medication delivery [33]. The CPPs' cationic charge density was a significant factor in determining the feasibility of the cargos' translocation. CPPs can be used to improve the specificity and selectivity of chemotherapy drugs by delivering them selectively to cancer cells while sparing normal cells. This approach not only reduces the side effects associated with traditional chemotherapy but also enhances the efficacy of the drugs by increasing their bioavailability and reducing the development of drug resistance. Extracellular vesicles altered with the hexadeca-arginine (R16) peptide revealed moderately good anti-cancer efficacy [85]. Proteins were typically transferred via CPPs by covalent bonding. Table 1 shows some of the CPPs used in the cancer treatment.

Unfortunately, because of changed biological activity and/or steric hindrance, covalent CPPs technology was not the most efficient method to transfer macromolecules [86]. Delivering oligonucleotides seems to be mostly accomplished by electrostatic adsorption. The linear structure of CPPs typically does not achieve highly satisfying oligonucleotide transfection efficiency due to low charge number, which led to poor complexation and organizational volatility of nano-carrier [33]. In order to create a unique bio-reducible cationic network employing R9 as a vector, Yoo et al., 2017 synthesised a branched R9 using disulfide bonds [87]. Branching structures enable effective electrostatic adsorption of pDNA or siRNA. In vitro, B-mR9 demonstrated strong intracellular trafficking and biocompatibility. The EPR impact of B-mR9 also demonstrated a targeted effect on the tumour that lasted for 48 hours. In the NCI-H460 harbouring BALB/c nude mice model, B-mR9/siVEGF significantly suppressed tumour development by 56.5% when compared to control, and the therapeutic effectiveness was greater than PEI25k and R9 vector [33].

A fresh method to create a platform for gene delivery was offered by the cationic network formed from CPPs. Moreover, ULK1 siRNA and the AMPK activator narciclasine might be delivered together in a pH-sensitive and biocompatible micelle system by Tai W., Gao X., 2017 in order to successfully prevent hepatocellular cancer in preclinical trials by controlling programmed cell death [86]. Subse-

quent studies confirmed CPPs were to be the revolutionary siRNA oligonucleotide paradigm. Clinical uses for CPPs were restricted to positive charges that caused systemic and off-target toxicity. In MCF-7 cells, CsA (Cyclosporin A) with electronic neutral, a new highly hydrophobic cyclic CPP, outperformed both pentapeptide VPT (VPTLQ) and PFV by a factor of several times and conventional neutral CPPs by a large margin [33]. By administering a membrane-impenetrable pro-apoptotic peptide (PAD), the effectiveness and toxicity of cyclosporin A were compared to TAT. The uptake of PDA was increased 2.2–4.7-fold in the tumour cell lines examined by CsA when CsA conjugated to PAD, and cellular uptake of CsA- PAD was often higher than TAT-PAD. Depending on the cell type, the cytotoxicity of CsA-PAD was comparable to or greater than TAT-PAD in four distinct tumour cell lines, but it was much more potent than PAD. CsA-PAD showed equivalent anti-tumor effect to TAT-PAD in xenografted MCF-7 nude mouse models, but with less systemic toxicity [33]. Although the correct tissue distribution of electro-neutral CPPs has to be further evaluated, cationic CPPs likely have more potential application value in vivo than electroneutral CPPs [88]. Hyaluronic acid (HA), a high-affinity ligand for tumour surface-specific overexpressed marker CD44, was one such polyanionic substance employed to coat nanoparticles in order to decrease toxicity and non-target of positive CPPs [33]. In order to administer 10-HCPT against hepatocellular carcinoma, Zhao et al., 2018 created a multifunctional liposome modified with TAT and HA (HA/CPPs-10-HCPT-NPs) [89]. Low-intensity focused ultrasound was employed to precisely regulate drug release to tumour tissue. In the multicellular tumour spheroid model, the liposome penetration depth enhanced 2.76- fold following TAT modification. HA-coated nano-carrier was a useful and promising method for CPPs administration in vivo [33]. Liposome coupled application of HA and CPPs with the assistance of ultrasound had a considerably greater tumour inhibition against hepatic carcinoma than other groups [89]. Figure 3 below shows the types of cancer which were tested with CPPs.




Figure 3: Therapeutic applications of CPPs in cancer treatment. This figure illustrates applications of CPPs in cancer therapy, centering on their role in protein mimicry to allow various therapeutic interventions. CPP has an enhanced chemotherapeutic efficacy as it improves intracellular drug delivery and pharmaco-

kinetics, and also allows the modulation of protein–protein interactions critical for disrupting oncogenic signaling pathways. CPPs also can act as vehicles for antibody delivery and immune detection, as well as for conjugation strategies such as linking with drugs like to promote targeted cytotoxicity. Figure adapted from <https://doi.org/10.1016/j.addr.2021.114044>

Poor prognosis, particularly for duodenal cancer, gliomas, and lymph metastases, was a significant obstacle in cancer treatment. Refractory cancers have a limited capacity to be treated since delivery methods couldn't get medications to the treatment site due of the complicated tumour microenvironment. CPPs may serve as the molecular motor for tumor-deepening cargo. To create mixed micelle for siRNA systemic administration, two tandem peptides (pTP-PEG- iRGD and pTP-iRGD) were synthesised by Lo et al., 2018 in an effort to solve the target and tumour stroma penetration problems in PDAC [90]. It could successfully get beyond PDAC's delivery obstacles to penetrate tumours in three-dimensional organoids and models of native tumours. Moreover, the mixed micelle complexed siRNA greatly slowed the development of the tumour [90]. For the purpose of treating gliomas, CPPs may cause cargos to pass the blood–brain barrier (BBB). To transport siRNA against glioma, Liu et al., 2018 created an anionic random- coiled polypeptide (PLG) coated CPPs (PVBLG-8) micelle [91]. To create a stable structure in serum and turn the micelle's surface potential negative, PLG became entangled with the PVBLG- 8/siRNA complex [91]. Also, micelle could carry out PVBLG-8's cell penetrating activity in response to low pH in the tumour extracellular microenvironment. CPPs were paired with glioma- homing peptides to precisely translocate siRNA in order to increase the target limit of CPPs in glioma applications [33, 91]. To maximise tumor-specific targeting and gene knockdown impact, the bonding form of two CPPs (PF14, PF28) with targeting moistures via either covalent conjugation or non-covalent complex was adjusted [33]. Lymph nodes nearby were the site of the first tumour metastasis and eventually extended across the entire body, making lymph metastasis an important channel of tumour spread [33]. Because of the blood-lymph barrier, current medicines that target lymphatic metastasis by intravenous injection have limited non-target and low penetration capability. After being administered intravenously for a possible anti-metastasis treatment, a R9 modified cabazitaxel nanoparticle (R9-CN) with a 13 nm size and a little positive charge was shown to have a conspicuous lymph target and deep penetration impact [33]. R9-CN's fluorescence signal persisted in primary tumour locations for at least 24 hours at a high intensity [92]. In a breast cancer lymphatic metastasis model, R9-CN significantly reduced the tumour growth rate by 1.4-fold and demonstrated a 63.3% inhibition rate of lung metastasis compared to CN. Deep lymphatic penetration made CPPs modified nanoparticles an excellent anti-metastasis platform [92].

Table 1: Current list of CPP for cancer

CPP	Type	Cancer Types	Function	Mechanism	Clinical Trial ID
PEP-010	Cationic	Breast cancer	Restores apoptotic pathways	Disrupts the caspase-9/PP2A interaction, activating caspase-dependent apoptosis	NCT04733027 (Phase I)
ATX-101	Cationic	Multiple myeloma/Sarcoma	Reaches the cell nucleus to enhance damage repair, cellular stress response, and the efficacy of several anti-cancer agents	Disrupts the PCNA/A PIM-containing protein interaction	NCT05116683 (Phase II), NCT04814875 (Phase I/II), NCT01462786 (Phase I)
AVB-620	Cationic	Breast cancer	Real-time tumour visualization during surgery	CPP conjugated with fluorophores Cy5/Cy7 for FRET, targeted to human breast cancer cells due to their MMP overexpression	NCT02391194 (Phase I), NCT03113825 (Phase II)

Z12 and ZEB-RA-Derived CPPs	Cationic	Broad spectrum, including aggressive brain cancers	Component of cancer vaccines	Promotes immune responses against tumors when conjugated with multi-epitopic antigens	NCT04046445 (Phase I)
pVEC and PEGA	Cationic	Breast cancer	Targeted drug delivery vector	Selective non-endocytic translocating mechanism (pVEC) by targeting molecular markers on tumor cells when conjugated with homing peptides (PEGA)	
Pep-1	Cationic	Broad spectrum	Targeted macromolecular carrier and drug delivery vector	High selective non-endocytic translocation through cancer cell membranes is primarily due to the high presence of acidic components	

MAP	Cati-onic	Broad spec-trum	Bifunc-tional CPP that dis-rupts cancer cell mem-branes	Selec-tive strong electro-static interac-tions with nega-tively charged phos-pholip-ids	
p28	Cati-onic	Multiple cancer types, includ-ing gli-oblas-toma and hepato-cellular carci-noma	Pro-motes cell-cycle arrest and apop-tosis in tumor cells	Interacts with wild-type and mutant p53 proteins, inhibit-ing their ubiquiti-nation and regulat-ing their levels	NCT0 09149 14 (Phas e I), NCT0 19751 16 (Phas e I), NCT0 53598 61 (Phas e II), NCT0 61025 25 (Phas e I)
SAP and SAP(E)	Pro-line-rich amphi-patic	Broad spec-trum	Tar-geted drug deliv-ery vector with mini-mal toxicity	Specific electro-static interac-tions with nega-tively charged mem-brane compo-nents (SAP); internal-ization of ag-gre-gates in a non-clathrin or cave-oline-mediat-ed endocyto-sis (SAP(E)	

				)	
Bac1-24	Pro-line-rich amphi-patic	Broad spec-trum, particu-larly solid tumors	Tar-geted deliv-ery agent of ther-a-peutic pro-teins and pep-tides	Hydro-phobic domains and specific electro-static interac-tions with nega-tively charged phos-pholip-ids	
BIM-SAHB _A	Sta-pled pep-tide	Hema-tologic cancers	Re-stores apop-tosis in re-sistant cancer cells	Blocks the anti-apoptot-ic se-questra-tion of BAX/BA K BH3 helices, mimick-ing the BH3 death domain	
SAHB _D	Sta-pled pep-tide	Can-cers where MCL-1 overex-pres-sion is a critical survival factor (mye-loma, acute myeloid leuke-mia, mela-noma, etc.)	Re-stores apop-tosis in re-sistant cancer cells	Inhibits the MCL-1 anti-apoptot-ic activi-ty, dis-rupting its inter-action with pro-apoptot-ic pro-teins	
ALRN-6924	Sta-pled pep-tide	Broad spec-trum, includ-ing breast cancer and acute	Re-stores p53 func-tion, reac-tivating apop-tosis	Binds strongly to MDM2 and MDMX, inhibit-ing the p53	NCT0 22646 13 (Phas e I/II), NCT0 40228 76 (Phas

		myeloid leukemia		suppression	e I), NCT03654716 (Phase I), NCT05622058 (Phase)
P1pal-7	Pepducin	Breast, lung, and ovarian cancer	Reduces tumor growth and slows cancer progression. Anti-angiogenic agent	Interacts with PAR1, inhibiting its activation	
EN1-iPeps	Homeodomain-derived	Breast cancer	Triggers a selective apoptosis response	Inhibits the EN1 transcription factor in tumor cells where it is over-expressed	
Vecto-cell@/DPVs	HS Binding CPP	Broad spectrum	Targeted drug delivery agent (from small compounds to macromolecules)	Caveolae-mediated endocytosis of DPVs-Glycosaminoglycan clusters	
CPPecp	HS Binding CPP	Tumours with high HS expression, including colon cancer	Inhibits cancer cell migration and angiogenesis	Binding to over-expressed heparan sulfate on the surface of cancer cells	

Melittin and derivatives	Derived from animal venoms and toxins	Broad spectrum	Drug delivery vector and apoptosis inducers in tumor-associated macrophages	An amphipathic α -helix structure enables interactions with the membrane, allowing the internalization of conjugated pro-apoptotic peptides	
Lycosin-I and R-Lycosin-I	Derived from animal venoms and toxins	Broad spectrum	Induces apoptosis in cancer cells and inhibits cell proliferation	Activates the mitochondrial death pathway and upregulates p27	
Pardaxins	Derived from animal venoms and toxins	Broad spectrum including aggressive cancers such as ovarian cancer and oral squamous cell carcinoma	Apoptosis inducer	Generation of ROS and mitochondrial membrane depolarization	
BT1718	Cyclic	Solid and refractory tumours	Selective release of cytotoxic agents	Binds to over-expressed MT1-MMP in tumors realising DM1, a cytotoxic payload	NCT03486730 (Phase I/II)
¹⁷⁷ Lu-	Cyclic	SSTR2-	Selec-	This	NCT0

DO-TA ⁰ -Tyr ³ -Oc-treatate (Luta-thera®)		positive neuro-endo-crine tumours	tive deliv-ery of cyto-toxic agent	radio-conju-gate utilizes the so-mato-statin analog TATE to target SSTR2-positive neuro-endo-crine tumors, deliver-ing a cytotox-ic dose of β radi-ation	2125474 (Phas e II), NCT02236910 (Phas e II), NCT03325816 (Phas e I/Phas e II)
---	--	-----------------------------------	------------------------------------	--	--

Adapted from: Moreno-Vargas, L. M., & Prada-Gracia, D. (2025). Cancer-Targeting Applications of Cell-Penetrating Peptides. *International Journal of Molecular Sciences*, 26(1), 2. <https://doi.org/10.3390/ijms26010002>

6. Use of CPPs in the Treatment of Inflammatory Conditions

The stratum corneum and mucosa are the primary delivery-related challenges with transdermal administration, which has high compliance and is an efficient method of local delivery of anti- inflammatory medicines. Polyargi-nine peptides are frequently used in transdermal medica-tion administration because of their effective ability to penetrate skin. To treat rat paw edoema, Gao et al., 2019 created lornoxicam-loaded lipid gels that were R11 mod-ified (LN-NLC-R11) [93]. Rat paw edoema was eliminat-ed, and LN-NLC-R11 dramatically reduced the generation of inflammatory cytokines as compared to NLC in vivo [93]. The distance between CPPs and nanoparticles will affect the effectiveness of cellular internalisation because of steric hindrance.

Due to steric hindrance, the separation between CPPs and nanoparticles will affect the effectiveness of cellular inter-nalisation. To reduce pulmonary inflammation, CPPs mod-ified gene carriers R9Gn-chitosan/siMIF (n = 0, 4, 10) were developed [33]. R9Gn-chitosan/in siMIF's vivo cell uptake, gene silencing effectiveness, and anti-inflammatory efficacy were all enhanced by lengthening the Gn controlled spacer arm [33]. In a mouse model of particulate matter-induced airway inflammation, R9G10-chitosan/siMIF dramatically decreased inflammation and goblet cell hyperplasia of lung tissue compared to R9-chitosan/siMIF [94]. Similar to this, intranasal injection of phospholipase D1 (PLD1) conjugated with TAT enhanced the anti-asthmatic action [95]. Epidermal hyperplasia is a

prevalent condition with a significant immune cell infiltra-tion known as psoriasis. A significant contributor to the pathophysiology of psoriasis is signal transducer and acti-vator of transcription 3 (STAT3). The high-affinity peptide designed to inhibit STAT3 is called APTstat3. R9 (APTstat3-9R) altered APTstat3 to increase stratum corneum penetration [33]. After topical intradermal thera-py, APTstat3-9R showed positive efficacy in reducing localised psoriasis-like skin irritation [33]. APTstat3-9R, however, had minimal skin penetration after transcutane-ous injection because the stratum corneum barrier inhibit-ed it. APTstat3-9R complexed with DMPC/DHPC to cre-ate discoidal lipid nanoparticles (DLNPs) with great col-loidal stability (20 nm size) and minimal side effects, im-proving transdermal administration [33]. In a mouse model of psoriasis, DLNPs were able to pass through the stratum corneum because of their lipophilicity and then pass through the openings between epidermal layers to reach the dermal layers [96]. The imiquimod-induced psoriatic mouse model's skin edoema and epidermal hyperplasia were successfully decreased by DLNPs [96].

Restrictions of the Use of CPPs

Being one of the most effective methods for transfection in many cell types, CPPs have been employed more and more over the past 30 years in the diagnosis and treatment of various diseases. In order to improve the treatment result, CPPs have proved essential in obtaining therapeutic con-centrations in cells and tissue that are challenging to target. Its success is based on both their flexibility and their po-tent transmembrane delivery feature [33]. They can easily be enhanced, changed, and synthesised.

CPPs are a two-edged sword, though, since they may have serious negative consequences for a number of reasons. FDA has not yet authorised any CPP-conjugated medica-tions, and numerous clinical trials have been stopped. One of the major limitations of CPPs is their poor specificity for target cells and tissues. CPPs can penetrate virtually any cell membrane, including healthy cells, which can result in off-target effects and unwanted toxicity. This lack of selectivity can be particularly problematic for cancer treatment, where the goal is to target cancer cells specifi-cally while sparing healthy cells. To overcome this limita-tion, several strategies have been developed to enhance the specificity of CPPs, such as conjugating them with target-ing moieties, such as antibodies or aptamers, that recog-nize cancer-specific markers or receptors. The quick con-sent from blood has a downside in that enzymatic break-down might cause a therapeutic payload to deteriorate in circulation before to reaching the therapeutic location. Because to its polypeptide nature, CPPs will enhance the likelihood that patients may have an unfavourable immu-nological reaction, which might both lower the efficacy of the medicine and result in an unpleasant immune stress response [97]. To increase exposure and decrease the im-munogenicity of treatments, it can be administered in high

doses and at regular intervals, however doing so causes toxicity. The alternative method is to provide treatments subcutaneously, which can lessen the immune system's tendency to produce anti-drug antibodies [98]. CPPs can be internalised by practically all types of cells due to treatments being absorbed by normal tissues off- target by cells. However, it should be noted that anything might become cytotoxic at a high dosage, and many concerns need to be solved before being formally administered to the patients [33]. CPPs are foreign molecules that can trigger immune responses, leading to inflammation and other adverse effects. They can accumulate in certain organs, such as the liver and kidney, which can cause toxicity and organ damage over time [33]. To address these issues, researchers have focused on developing CPPs with reduced immunogenicity and toxicity profiles, as well as improving their pharmacokinetic properties to enhance their clearance from the body. The majority of study revealed minimal toxicity of CPPs. CPPs' cytotoxicity is a major basis of apprehension [99]. As a typical CPP, the MAP, for instance, shares structural similarities with antimicrobial lytic peptides that impact microorganisms by disrupting their plasma membranes [100]. In addition to these limitations, CPPs also face several challenges in molecular diagnostics and cancer treatment. One of the challenges is the efficient delivery of CPP-cargo conjugates to their intended targets. CPPs are often used to deliver cargoes, such as nucleic acids, to specific cell types, such as cancer cells, where they can regulate gene expression or induce cell death. However, efficient delivery of these conjugates to their intended targets can be challenging due to the complex and heterogeneous nature of tumors and the extracellular matrix that surrounds them. Another challenge is the development of CPP-based therapeutics that can overcome drug resistance mechanisms. Cancer cells can develop resistance to chemotherapy drugs and other targeted therapies, which can limit their effectiveness. To address this issue, researchers have focused on developing CPP-based therapeutics that can bypass drug resistance mechanisms and enhance the efficacy of existing therapies. MAPs can cause the leaking of protons, proteins, metal ions, etc. due to their amphipathic action in the presence of artificial micelles, which leads to cell death from plasma membrane injury [33, 101]. The ability of cationic CPPs to attach to glycosaminoglycans is well acknowledged, however it is unclear if CPPs can interact with certain membrane receptors [102]. Because of a reduced local concentration, the efficacy of CPP-conjugated therapies can be decreased in tissues with extensive tissue distribution [33]. Consequently, it's critical to target certain cells as much as possible while disregarding normal cells. Another major issue is the transport of CPPs and CPP/cargo complexes to the cytosol from endosomes prior to lysosomal destruction [103]. It is thought that medications that remain inside endosomes are unable to perform biologically. In order to hasten the release of the carrier from the endosome into the cytosol, CPPs should be designed in a way that efficiently promotes endosomal escape [33]. To achieve effective thera-

py, the most crucial ways for CPP delivery are not only translocated to the target tissue or organ but also target into certain organelles in the cell, such as the nucleus and [104]. To summarise, it is needed urgently to develop optimal CPPs with low toxicity, high efficiency, and specificity in order to fulfil the clinical use of CPPs and to solve therapeutic problems related to CPPs. Despite these challenges and limitations, CPPs have the potential to revolutionize molecular diagnostics and cancer treatment. Their ability to penetrate cell membranes and deliver various types of cargoes to cells has opened up new avenues for the development of targeted therapies and precision medicine. In the future, further research is needed to address the challenges and limitations of CPPs and to fully realize their clinical potential.

Future Prospects

In clinical trials, evaluating the efficacy of a therapeutic agent is paramount; however, several other critical factors must be considered for successful drug development. These factors include large-scale production feasibility, physicochemical properties, stability, toxicity, pharmacokinetics, formulation strategies, and the chosen route of administration. For CPPs, despite their ability to efficiently transport biomolecules across cellular membranes, challenges such as poor specificity, limited stability, and rapid clearance from the body restrict their clinical applicability [105]. To enhance their pharmacological performance, multiple strategies have been explored to address these limitations.

Cationic CPPs have demonstrated an ability to improve the intracellular delivery of large hydrophilic biomolecules, including peptide nucleic acids and oligonucleotides. However, their uptake primarily occurs via endocytic pathways, necessitating effective mechanisms to promote endosomal escape to maximize bioavailability. Promising approaches involves conjugating lipid moieties, particularly fatty acids, to cationic peptides, which can enhance their biological activity. The efficacy of these lipid modifications depends on the length of the fatty acid chain, with longer chains (ranging from C8 to C16) showing progressively increased effects [105]. However, this enhancement comes at a cost, as cytotoxicity also rises with increasing fatty acid chain length.

Extensive in vitro investigations have indicated that CPPs generally exhibit low toxicity and minimal immunogenicity. However, their immunogenic potential varies depending on physicochemical characteristics such as molecular size, charge distribution, amino acid composition, structural morphology, and the specific type of conjugated cargo [105]. These attributes collectively influence both membrane integrity and the probability of eliciting an immune response. Consequently more studies are required to fully understand and mitigate any immunological risks associat-

ed with CPPs. The fusion or conjugation of CPPs with therapeutic cargoes, such as drugs or vaccines, has been shown to mitigate their inherent toxicity. Nonetheless, precise dose optimization remains a crucial aspect of research in the development of CPP-based therapeutics.

One of the most significant barriers to the clinical application of CPPs is their vulnerability to enzymatic degradation in biological environments. Upon exposure to serum or cellular conditions, CPPs are readily degraded by proteases present in bodily fluids such as blood, gastric and intestinal secretions, extracellular matrix, and intracellular compartments. This rapid degradation severely limits their systemic half-life, thereby reducing therapeutic efficacy. The reticuloendothelial system, particularly in the liver and spleen, actively eliminates positively charged molecules, further restricting the circulation time of peptide-based therapeutics [105]. Addressing these challenges requires strategic chemical modifications that hinder proteolytic recognition and degradation, thereby prolonging the half-life and enhancing cellular uptake.

Traditional approaches to improve peptide stability include modifications at the peptide termini. N-terminal acetylation and C-terminal amidation are commonly employed to shield the peptide backbone from enzymatic cleavage. A more advanced and effective strategy involves stereochemical modifications, such as incorporating D-amino acids or non-natural amino acids [105]. These modifications disrupt enzymatic recognition, significantly extending the intracellular retention of CPPs. However, while such alterations can enhance proteolytic resistance, excessive modifications can introduce undesirable toxic effects. For instance, a study comparing two CPP derivatives found that while one retained low toxicity, a fully modified counterpart containing only D-amino acids exhibited severe toxicity *in vivo*.

An alternative approach to improve CPP stability involves the physical shielding of peptides using hydrophilic polymers, such as polyethylene glycol (PEG). PEGylation can protect CPPs from enzymatic degradation, enhance their metabolic stability, prolong circulation time, and reduce immunogenicity [105]. Despite these advantages, PEGylation also presents a major drawback: it diminishes CPP interactions with cell membranes, ultimately reducing cellular uptake efficiency. To overcome this limitation, precise control over polymer size and conjugation density is required. Researchers have thus explored strategies to facilitate the detachment of PEG near the target site to restore CPP functionality. However, concerns regarding PEG stability, particularly its susceptibility to oxidative degradation and limited excretion, have prompted investigations into alternative polymers. Polyglycerols, including linear, dendritic, and hyperbranched variants, have emerged as promising substitutes due to their superior resistance to oxidation and thermal stress.

Comparative studies have shown that cyclic CPPs exhibit superior internalization efficiency and stability over their linear counterparts. This enhanced performance is attributed to their increased affinity for target receptors, which facilitates more effective cellular entry. Unlike linear peptides, cyclic CPPs display greater rigidity, reducing entropy loss and allowing tighter interactions with phospholipids in fluid membranes [105]. This property is particularly advantageous for endosomal escape, as higher membrane affinity leads to more efficient transduction. Research has further demonstrated that specific modifications, such as peptide suturing via covalent cross-linking of amino acid side chains, can significantly enhance metabolic stability and membrane permeability by stabilizing the peptide's conformation.

Conclusion

Various biomacromolecules, including polypeptides, proteins, and nucleic acids, can be transported into cells by CPPs through the cell membrane, which transcends the limitations of conventional drug administration, and opens up new possibilities for the study of drug carriers, and producing positive outcomes. There has been advancement, particularly in the areas of tumour therapy, cardiovascular illness, nervous system disease, and vaccine research, but a number of issues still need to be resolved before CPPs can truly be used in clinical practise. One is that proteolysis and quick liver and kidney clearance cause CPPs to have a short plasma half-life when transporting proteins or peptides, which leads to poor CPP stabilization. It may be essential to double-double embed CPPs and medications into macromolecular carriers (such as liposomes or biopolymers) in order to solve this issue. Another option is to link hydrophilic PEG. The second is the lack of tissue and cell type specificity when CPPs facilitate transmembrane, which prevents the drug from building up in the target site and results in relatively significant damage to healthy tissues and cells. To get around this issue, several researchers have attempted to construct a number of modified CPPs that cover up their membrane-penetrating properties before reaching the target site and take advantage of alterations in the tumour microenvironment once they have arrived there. In order to achieve anti-tumor targeted therapy, CPPs' transmembrane transport function can be restored through, for instance, changes in pH value, enzyme type and activity, or the application of external stimuli. The study of the CPPs delivery system is currently becoming increasingly in-depth. A new concept for the entry of exogenous biomolecules into cells is presented by the transduction technology facilitated by cell-penetrating peptides, which offers promise for the treatment of disorders. The number of cell-penetrating peptides and their analogues will increase as the research goes on, and their analogues' membrane-penetrating mechanisms will be further investigated.

List of Abbreviations

CPP Cell-Penetrating Peptides	MPG-8 (A shortened version of MPG; specific peptide variant)
TAT Trans-activator of Transcription	CLIO Cross-Linked Iron Oxide particles
BBB Blood-Brain Barrier	MRI Magnetic Resonance Imaging
HIV Human Immunodeficiency Virus	HA Hyaluronic Acid
PTDs protein transduction domains	10-HCPT 10-Hydroxycamptothecin
NLS Nuclear Localization Signal	NPs Nanoparticles
SV40 Simian Virus 40	PDAC Pancreatic Ductal Adenocarcinoma
NMR Nuclear Magnetic Resonance	PLG (An anionic random-coiled polypeptide; designated as PLG)
PPII Polyproline II	PVBLG-8 (A specific cell-penetrating peptide)
FGF Fibroblast Growth Factor	PF14 (A specific CPP variant)
CME Clathrin-Mediated Endocytosis	PF28 (A specific CPP variant)
CvME Caveolin-Mediated Endocytosis	CsA Cyclosporin A
AP-2 Adaptor Protein Complex 2	PAD Pro-Apoptotic Peptide
FM4-64 (a fluorescent endocytic marker; typically referred to by its trade name)	EPR Enhanced Permeability and Retention
CHO Chinese Hamster Ovary	pDNA Plasmid DNA
HEK293 Human Embryonic Kidney 293	AMPK AMP-Activated Protein Kinase
MDCK Madin-Darby Canine Kidney	ULK1 Unc-51 Like Autophagy Activating Kinase 1
TP Transportan	iRGD Internalizing RGD (Arg-Gly-Asp) peptide
125I Iodine-125	PEG Polyethylene Glycol
68Ga Gallium-68	R9-CN R9-modified Cabazitaxel Nanoparticle
64Cu Copper-64	CN Cabazitaxel Nanoparticle
99mTc Technetium-99m	R16 (Hexadeca-Arginine; 16 arginine residues)
111In Indium-111	MCF-7 Michigan Cancer Foundation-7 (breast cancer cell line)
177Lu Lutetium-177	R11 An 11-mer arginine peptide used for modification
GFP Green Fluorescent Protein	LN-NLC-R11 Lornoxicam-Loaded Nanostructured Lipid Carrier modified with R11
GAGs Glycosaminoglycans	R9Gn-chitosan/siMIF A gene carrier composed of R9Gn-chitosan complexed with siRNA targeting MIF (Macrophage Migration Inhibitory Factor)
F-actin Filamentous Actin	R9G10-chitosan/siMIF A variant of the gene carrier where the spacer consists of 10 glycine residues (R9G10) complexed with siRNA targeting MIF
FAM Fluorescein Isothiocyanate	PLD1 Phospholipase D1
TAMRA Tetramethylrhodamine	STAT3 Signal Transducer and Activator of Transcription 3
MDR Multidrug Resistance	APTstat3-9R A high-affinity peptide designed to inhibit STAT3 that is modified with 9 arginine residues
R8 (Arginine) Octamer (8 arginine residues)	DMPC 1,2-Dimyristoyl-sn-glycero-3-phosphocholine
R8-taxol A conjugate of R8 with Taxol (paclitaxel)	
R9 (Arginine) Nonamer (9 arginine residues)	
p53 Tumor Protein 53	
p16 p16 Protein (a cyclin-dependent kinase inhibitor)	
SMAC Second Mitochondrial-Derived Activator of Caspase	
TRAIL TNF-Related Apoptosis-Inducing Ligand	
RIP Ribosome Inactivating Protein	
LMWP Low Molecular Weight Protamine	
siRNA Small Interfering RNA	
VEGF Vascular Endothelial Growth Factor	

DHPC Dihexanoylphosphatidylcholine
DLNPs Discoidal Lipid Nanoparticles
FDA U.S. Food and Drug Administration
MAP Model Amphipathic Peptide

Acknowledgement

The author would like to acknowledge the use of Grammarly for proofreading and QuillBot for writing support, which helped enhance the clarity and coherence of the text.

Author Contribution Statement

The author is the sole contributor and takes full responsibility for the preparation, finalization, and approval of the manuscript.

Conflicts of Interest

Declared none.

Funding

No external funding was received for this research.

References

- [1] Green, M. and Loewenstein, PM (1988) Autonomous Functional Domains of Chemically Synthesized Human Immunodeficiency Virus Tat Trans-Activator Protein. *Cell*, 55, 1179-1188. [https://doi.org/10.1016/0092-8674\(88\)90262-0](https://doi.org/10.1016/0092-8674(88)90262-0)
- [2] Chen Yueru, Wu Mengqian, Hu Qian, Guo Huarong. Research overview of cell-penetrating peptides [J]. *Frontiers of Marine Science*, 2018, 5(2): 29-37. <https://doi.org/10.12677/AMS.2018.52004>
- [3] Claudio Brigati, Mauro Giacca, Douglas M. Noonan, Adriana Albini, HIV Tat, its TARgets and the control of viral gene expression, *FEMS Microbiology Letters*, Volume 220, Issue 1, March 2003, Pages 57–65, [https://doi.org/10.1016/S0378-1097\(03\)00067-3](https://doi.org/10.1016/S0378-1097(03)00067-3)
- [4] Milletti, F. (2012) Cell-Penetrating Peptides Classes, Origin and Current Landscape. *Drug Discovery Today*, 17, 850-860. <https://doi.org/10.1016/j.drudis.2012.03.002>
- [5] Xu J., Khan A. R., Fu M., Wang R., Ji J., Zhai G. (2019). Cell-penetrating peptide: a means of breaking through the physiological barriers of different tissues and organs. *J. Control Release* 309, 106–124. <https://doi.org/10.1016/j.jconrel.2019.07.020>
- [6] Skotland, T., Iversen, T. G., Torgersen, M. L., & Sandvig, K. (2015). Cell-penetrating peptides: possibilities and challenges for drug delivery in vitro and in vivo. *Molecules* (Basel, Switzerland), 20(7), 13313–13323. <https://doi.org/10.3390/molecules200713313>
- [7] Guo, Z., Peng, H., Kang, J., & Sun, D. (2016). Cell-penetrating peptides: Possible transduction mechanisms and therapeutic applications. *Biomedical reports*, 4(5), 528–534. <https://doi.org/10.3892/br.2016.639>
- [8] Chu D., Xu W., Pan R., Ding Y., Sui W., Chen P. (2015). Rational modification of oligoarginine for highly efficient siRNA delivery: structure–activity relationship and mechanism of intracellular trafficking of siRNA. *Nanotechnol. Biol. Med.* 11,

435–446. 10.1016/j.nano.2014.08.007

DOI:
10.1080/17425247.2019.1676720

[9] Tunnemann, G., Ter-Avetisyan, G., Martin, RM, et al. (2008) Live-Cell Analysis of Cell Penetration Ability and Toxicity of Oligo-Arginines. *Journal of Peptide Science*, 14, 469-476.

<https://doi.org/10.1002/psc.968>

[10] Futaki, S., Suzuki, T., Ohashi, W., et al. (2001) Arginine-Rich Peptides. An Abundant Source of Membrane Permeable Peptides Having Potential as Carriers for Intracellular Protein Delivery. *The Journal of Biological Chemistry*, 276, 5836-5840.

<https://doi.org/10.1074/jbc.M007540200>

[11] Prochiantz, A. (1996) Getting Hydrophilic Compounds into Cells: Lessons from Homeopeptides. *Current Opinion in Neurobiology*, 6, 629-634.

[https://doi.org/10.1016/S0959-4388\(96\)80095-X](https://doi.org/10.1016/S0959-4388(96)80095-X)

[12] Kimia Kardani, Alireza Milani, Samaneh H. Shabani & Azam Bolhassani (2019) Cell penetrating peptides: the potent multi-cargo intracellular carriers, *Expert Opinion on Drug Delivery*, 16:11, 1227-1258,

[13] Eiríksdóttir E., Konate K., Langel U., Divita G., Deshayes S. (2010). Secondary structure of cell-penetrating peptides controls membrane interaction and insertion. *Biochim. Biophys. Acta* 1798, 1119–1128. 10.1016/j.bbame.2010.03.005

[14] Milletti F. (2012). Cell-penetrating peptides: classes, origin, and current landscape. *Drug Discovery Today* 17, 850–860. 10.1016/j.drudis.2012.03.002

[15] Oehlke, J., Krause, E., Wiesner, B., et al. (1997) Extensive Cellular Uptake into Endothelial Cells of an Amphipathic β -Sheet Forming Peptide. *FEBS Letters*, 415, 196-199. [https://doi.org/10.1016/S0014-5793\(97\)01123-X](https://doi.org/10.1016/S0014-5793(97)01123-X)

[16] Carnevale K. J. F., Muroski M. E., Vakil P. N., Foley M. E., Laufer G., Kenworthy R., et al. (2018). Selective Uptake Into Drug Resistant Mammalian Cancer by Cell Penetrating Peptide-Mediated Delivery.

Bioconjugate Chem. 29, 3273–3284. 10.1021/acs.bioconjchem.8b00429

[17] Kauffman W. B., Fuselier T., He J., Wimley W. C. (2015). Mechanism Matters: A Taxonomy of Cell Penetrating Peptides. *Trends Biochem. Sci.* 40, 749–764. 10.1016/j.tibs.2015.10.004

[18] Borrelli A., Tornesello A., Tornesello M., Buonaguro F. (2018). Cell Penetrating Peptides as Molecular Carriers for Anti-Cancer Agents. *Molecules* 23, 295. 10.3390/molecules23020295

[19] Tashima T. (2017). Intelligent substance delivery into cells using cell-penetrating peptides. *Bioorg. Med. Chem. Lett.* 27, 121–130. 10.1016/j.bmcl.2016.11.083

[20] Brock, R. (2014) The Uptake of Arginine-Rich Cell-Penetrating Peptides: Putting the Puzzle Together.

Bioconjugate Chemistry, 25, 863-868. <https://doi.org/10.1021/bc500017t>

[21] Islam M. Z., Sharmin S., Moniruzzaman M., Yamazaki M. (2018). Elementary processes for the entry of cell-penetrating peptides into lipid bilayer vesicles and bacterial cells. *Appl. Microbiol. Biotechnol.* 102, 3879–3892. 10.1007/s00253-018-8889-5

[22] Guidotti G., Brambilla L., Rossi D. (2017). Cell-Penetrating Peptides: From Basic Research to Clinics.

Trends Pharmacol. Sci. 38, 406–424. 10.1016/j.tips.2017.01.003

[23] Munjal A., Khandia R., Kumar A., Singh G., Karthik K.,

Dhama K. (2017). Cell Penetrating Peptides: Biomedical/Therapeutic Applications with Emphasis as Promising Futuristic Hope for Treating Cancer. *Int. J. Pharmacol.* 13, 677–689. 10.3923/ijp.2017.677.689

[24] Shin, MC, Zhang, J., Min, KA, et al. (2014) Cell-Penetrating Peptides: Achievements and Challenges in Application for Cancer Treatment. *Journal of Biomedical Materials Research Part A*, 102, 575-587. <https://doi.org/10.1002/jbm.a.34859>

[25] Zhu P., Jin L. (2017). Cell Penetrating Peptides: A Promising Tool for the Cellular Uptake of Macromolecular Drugs. *Curr. Protein Pept. Sci.* 19, 211–220. 10.2174/1389203718666170710115240

[26] Pouny, Y., Rapaport, D., Mor, A., et al. (1992) Interaction of Antimicrobial Dermaseptin and Its Fluorescently Labeled Analogues with Phospholipid Membranes. *Biochemistry*, 31, 12416-12423. <https://doi.org/10.1021/bi00164a017>

[27] Matsuzaki, K. (1999) Why and How Are Peptide-Lipid Interactions Utilized for Self-Defense? Magainins and Tachyplesins as Archetypes. *Biochimica et Biophysica Acta (BBA)—Biomembranes*, 1462, 1-

10. [https://doi.org/10.1016/S0005-2736\(99\)00197-2](https://doi.org/10.1016/S0005-2736(99)00197-2)

- [28] Ludtke, S., He, K. and Huang, H. (1995) Membrane Thinning Caused by Magainin 2. *Biochemistry*, 34, 16764-16769. <https://doi.org/10.1021/bi00051a026>
- [29] Palm-Apergi, C., Lorents, A., Padari, K., et al. (2009) The Membrane Repair Response Masks Membrane Disturbances Caused by Cell-Penetrating Peptide Uptake. *FASEB Journal*, 23, 214-223. <https://doi.org/10.1096/fj.08-110254>
- [30] Yang Z., Xie J., Zhu J., Kang C., Chiang C., Wang X., et al. (2016). Functional exosome-mimic for delivery of siRNA to cancer: in vitro and in vivo evaluation. *J. Controlled Release* 243, 160–171. [10.1016/j.jconrel.2016.10.008](https://doi.org/10.1016/j.jconrel.2016.10.008)
- [31] Fittipaldi A., Ferrari A., Zoppé M., Arcangeli C., Pellegrini V., Beltram F., et al. (2003). Cell Membrane Lipid Rafts Mediate Caveolar Endocytosis of HIV-1 Tat Fusion Proteins. *J. Biol. Chem.* 278, 34141–34149. [10.1074/jbc.M303045200](https://doi.org/10.1074/jbc.M303045200)
- [32] Wadia J. S., Stan R. V., Dowdy S. F. (2004). Transducible TAT-HA fusogenic peptide enhances escape of TAT-fusion proteins after lipid raft macropinocytosis. *Nat. Med.* 10, 310–315. [10.1038/nm996](https://doi.org/10.1038/nm996)
- [33] Xie, J., Bi, Y., Zhang, H., Dong, S., Teng, L., Lee, R. J., & Yang, Z. (2020). Cell-Penetrating Peptides in Diagnosis and Treatment of Human Diseases: From Preclinical Research to Clinical Application. *Frontiers in pharmacology*, 11, 697. <https://doi.org/10.3389/fphar.2020.00697>
- [34] Vives, E., Granier, C., Prevot, P., et al. (1997) Structure-Activity Relationship Study of the Plasma Membrane Translocating Potential of a Short Peptide from HIV-1 Tat Protein. *Letters in Peptide Science*, 4, 429-436. <https://doi.org/10.1023/A:1008850300184>
- [35] Lundberg, M. and Johansson, M. (2002) Positively Charged DNA-Binding Proteins Cause Apparent Cell Membrane Translocation. *Biochemical and Biophysical Research Communications*, 291, 367-371. <https://doi.org/10.1006/bbrc.2002.6450>
- [36] Richard, JP, Melikov, K., Vives, E., et al. (2003) Cellpenetrating Peptides. A Reevaluation of the Mechanism of Cellular Uptake. *The Journal of Biological Chemistry*, 278, 585-590. <https://doi.org/10.1074/jbc.M209548200>
- [37] Patel, L. N., Zaro, J. L., & Shen, W. C. (2007). Cell penetrating peptides: intracellular pathways and

- pharmaceutical perspectives. *Pharmaceutical research*, 24(11), 1977–1992. <https://doi.org/10.1007/s11095-007-9303-7>
- [38] Conner, SD and Schmid, SL (2003) Regulated Portals of Entry into the Cell. *Nature*, 422, 37–44. <https://doi.org/10.1038/nature01451>
- [39] Pucadyil, T. J., & Schmid, S. L. (2009). Conserved functions of membrane active GTPases in coated vesicle formation. *Science (New York, N.Y.)*, 325(5945), 1217–1220. <https://doi.org/10.1126/science.1171004>
- [40] Aderem A., Underhill D. M. (1999). Mechanisms Of Phagocytosis In Macrophages. *Annu. Rev. Immunol.* 17, 593–623. [10.1146/annurev.immunol.17.1.593](https://doi.org/10.1146/annurev.immunol.17.1.593)
- [41] Zhao X.-L., Chen B.-C., Han J.-C., Wei L., Pan X.-B. (2015). Delivery of cell-penetrating peptide-peptide nucleic acid conjugates by assembly on an oligonucleotide scaffold. *Sci. Rep.* 5, 17640. [10.1038/srep17640](https://doi.org/10.1038/srep17640)
- [42] Ciobanasu, C., & Kubitcheck, U. (2020). Cell-Penetrating Peptides Targeting and Distorting Biological Membranes. *Surface and Interface Science: Volume 7: Liquid and Biological Interfaces*, 7, 441–469.
- [43] Xu J., Khan A. R., Fu M., Wang R., Ji J., Zhai G. (2019). Cell-penetrating peptide: a means of breaking through the physiological barriers of different tissues and organs. *J. Control Release* 309, 106–124. [10.1016/j.jconrel.2019.07.020](https://doi.org/10.1016/j.jconrel.2019.07.020)
- [44] Pujals S., Giralt E. (2008). Proline-rich, amphipathic cell-penetrating peptides. *Adv. Drug Deliv. Rev.* 60, 473–484. [10.1016/j.addr.2007.09.012](https://doi.org/10.1016/j.addr.2007.09.012)
- [45] McClorey G., Banerjee S. (2018). Cell-Penetrating Peptides to Enhance Delivery of Oligonucleotide-Based Therapeutics. *Biomedicines* 6, 51. [10.3390/biomedicines6020051](https://doi.org/10.3390/biomedicines6020051)
- [46] Ruseska, I.; Zimmer, A. Beilstein J. Nanotechnol. 2020, 11, 101–123. [doi:10.3762/bjnano.11.10](https://doi.org/10.3762/bjnano.11.10)
- [47] Madani, F., Lindberg, S., Langel, U., Futaki, S., & Gräslund, A. (2011). Mechanisms of cellular uptake of cell-penetrating peptides. *Journal of biophysics (Hindawi Publishing Corporation : Online)*, 2011, 414729. <https://doi.org/10.1155/2011/414729>
- [48] Fretz, M. M., Penning, N. A., Al-Taei, S., Futaki, S., Takeuchi, T., Nakase, I., Storm, G., & Jones, A. T. (2007). Temperature- and cholesterol-dependent translocation of L- and D-octa-arginine across the plasma and nuclear membrane of CD34+ leukaemia cells. *The Biochemical journal*, 403(2), 335–

342.
<https://doi.org/10.1042/BJ20061808>

0854.2007.00572.x

[49] Tünnemann, G., Martin, R. M., Haupt, S., Patsch, C., Edenhofer, F., & Cardoso, M. C. (2006). Cargo-dependent mode of uptake and bioavailability of TAT-containing proteins and peptides in living cells. *FASEB journal : official publication of the Federation of American Societies for Experimental Biology*, 20(11), 1775–1784. <https://doi.org/10.1096/fj.05-5523com>

[50] Maiolo, J. R., Ferrer, M., & Ottinger, E. A. (2005). Effects of cargo molecules on the cellular uptake of arginine-rich cell-penetrating peptides. *Biochimica et biophysica acta*, 1712(2), 161–172. <https://doi.org/10.1016/j.bbamem.2005.04.010>

[51] Jones, A. T., & Sayers, E. J. (2012). Cell entry of cell penetrating peptides: tales of tails wagging dogs. *Journal of controlled release : official journal of the Controlled Release Society*, 161(2), 582–591. <https://doi.org/10.1016/j.jconrel.2012.04.003>

[52] Duchardt, F., Fotin-Mleczek, M., Schwarz, H., Fischer, R., & Brock, R. (2007). A comprehensive model for the cellular uptake of cationic cell-penetrating peptides. *Traffic (Copenhagen, Denmark)*, 8(7), 848–866. <https://doi.org/10.1111/j.1600->

[53] Hällbrink, M., Oehlke, J., Papsdorf, G., & Bienert, M. (2004). Uptake of cell-penetrating peptides is dependent on peptide-to-cell ratio rather than on peptide concentration. *Biochimica et biophysica acta*, 1667(2), 222–228. <https://doi.org/10.1016/j.bbamem.2004.10.009>

[54] Mueller, J., Kretzschmar, I., Volkmer, R., & Boisguerin, P. (2008). Comparison of cellular uptake using 22 CPPs in 4 different cell lines. *Bioconjugate chemistry*, 19(12), 2363–2374. <https://doi.org/10.1021/bc800194e>

[55] Gluzman Y. (1981). SV40-transformed simian cells support the replication of early SV40 mutants. *Cell*, 23(1), 175–182. [https://doi.org/10.1016/0092-8674\(81\)90282-8](https://doi.org/10.1016/0092-8674(81)90282-8)

[56] Gronewold, A., Horn, M., Randelović, I., Tóvári, J., Muñoz Vázquez, S., Schomäcker, K., & Neundorff,

I. (2017). Characterization of a Cell-Penetrating Peptide with Potential Anticancer Activity. *ChemMedChem*, 12(1), 42–49. <https://doi.org/10.1002/cmdc.201600498>

[57] Zorko, M., & Langel, Ü. (2022). Studies of cell-penetrating peptides by biophysical methods.

Quarterly Reviews of Biophysics, 55,
E3. doi:10.1017/S0033583522000026

[58] Koo, J. H., Yoon, H., Kim, W. J., Lim, S., Park, H. J., & Choi, J. M. (2014). Cell membrane penetrating function of the nuclear localization sequence in human cytokine IL-1 α . *Molecular biology reports*, 41(12), 8117–8126. <https://doi.org/10.1007/s11033-014-3711-7>

[59] Wu, X., & Gehring, W. (2014). Cellular uptake of the Antennapedia homeodomain polypeptide by macropinocytosis. *Biochemical and biophysical research communications*, 443(4), 1136–1140. <https://doi.org/10.1016/j.bbrc.2013.12.062>

[60] He, J., Kauffman, W. B., Fuselier, T., Naveen, S. K., Voss, T. G., Hristova, K., & Wimley, W. C. (2013). Direct cytosolic delivery of polar cargo to cells by spontaneous membrane-translocating peptides. *The Journal of biological chemistry*, 288(41), 29974–29986. <https://doi.org/10.1074/jbc.M113.488312>

[61] Kim, HY, Yum, SY, Jang, G and Ahn, D-R (2015) Discovery of non-cationic cell penetrating peptide

derived from membrane-interacting human proteins and its potential as a protein delivery carrier. *Scientific Reports* 5, 11719.

[62] Ben-Dov, N., & Korenstein, R. (2015). The uptake of HIV Tat peptide proceeds via two pathways which differ from macropinocytosis. *Biochimica et biophysica acta*, 1848(3), 869–877. <https://doi.org/10.1016/j.bbamem.2014.12.015>

[63] Pooga, M., Hällbrink, M., Zorko, M., & Langel, U. (1998). Cell penetration by transportan. *FASEB journal : official publication of the Federation of American Societies for Experimental Biology*, 12(1), 67–77. <https://doi.org/10.1096/fasebj.12.1.67>

[64] Zaro, J. L., Vekich, J. E., Tran, T., & Shen, W. C. (2009). Nuclear localization of cell-penetrating peptides is dependent on endocytosis rather than cytosolic delivery in CHO cells. *Molecular pharmaceutics*, 6(2), 337–344. <https://doi.org/10.1021/mp800239p>

[65] Yeh, T. H., Chen, Y. R., Chen, S. Y., Shen, W. C., Ann, D. K., Zaro, J. L., & Shen, L. J. (2016).

Selective Intracellular Delivery of Recombinant Arginine Deiminase (ADI) Using pH-Sensitive Cell Penetrating Peptides To Overcome ADI Resistance in Hypoxic Breast Cancer Cells. *Mo-*

lecular pharmaceuticals, 13(1), 262–271.
<https://doi.org/10.1021/acs.molpharmaceut.5b00706>

[66] Velikyan I. (2015). 68Ga-Based radiopharmaceuticals: production and application relationship.

Molecules (Basel, Switzerland), 20(7), 12913–12943.
<https://doi.org/10.3390/molecules200712913>

[67] Bergmann, R., Splith, K., Pietzsch, J., Bachmann, M., & Neundorff, I. (2017). Biological characterization of novel nitroimidazole-peptide conjugates in vitro and in vivo. *Journal of peptide science : an official publication of the European Peptide Society*, 23(7-8), 597–609.
<https://doi.org/10.1002/psc.2995>

[68] Abbasi Gharibkandi, N., Conlon, J. M., & Hosseinimehr, S. J. (2020). Strategies for improving stability and pharmacokinetic characteristics of radiolabeled peptides for imaging and therapy. *Peptides*, 133, 170385.
<https://doi.org/10.1016/j.peptides.2020.170385>

[69] Dubikovskaya, EA, Thorne, SH, Pillow, TH, et al. (2008) Overcoming Multidrug Resistance of Small-Molecule Therapeutics through Conjugation with Releasable Octaarginine Transporters. *Proceedings of the National Academy of Sciences of the United States of America*, 105, 12–12133.
<https://doi.org/10.1073/pnas.0805374105>

[70] Lee, JY, Choi, YS, Suh, JS, et al. (2011) Cell-Penetrating Chitosan/Doxorubicin/TAT Conjugates for Efficient Cancer Therapy. *International Journal of Cancer*, 128, 2470–2480.
<https://doi.org/10.1002/ijc.25578>

[71] Hollstein, M., Sidransky, D., Vogelstein, B., et al. (1991) p53 Mutations in Human Cancers. *Science*, 253, 49–53.
<https://doi.org/10.1126/science.1905840>

[72] Liggett, WH and Sidransky, D. (1998) Role of the p16 Tumor Suppressor Gene in Cancer. *Journal of Clinical Oncology*, 16, 1197–1206.

<https://doi.org/10.1200/JCO.1998.16.3.1197>

[73] Snyder, EL, Meade, BR, Saenz, CC, et al. (2004) Treatment of Terminal Peritoneal Carcinomatosis by a Transducible p53-Activating Peptide. *PLOS Biology*, 2, E36.

<https://doi.org/10.1371/journal.pbio.0020036>

[74] Hosotani, R., Miyamoto, Y., Fujimoto, K., et al. (2002) Trojan p16 Peptide Suppresses Pancreatic Cancer Growth and Prolongs Survival in Mice. *Clinical Cancer Research*, 8,

1271-1276.

[75] Vucic, D., Deshayes, K., Ackerly, H., et al. (2002) Smac Negatively Regulates the Anti-Apoptotic Activity of Melanoma Inhibitor of Apoptosis (MLIAP). *The Journal of Biological Chemistry*, 277, 12275-12279.

<https://doi.org/10.1074/jbc.M112045200>

[76] Fulda, S., Wick, W., Weller, M., et al. (2002) Smac Agonists Sensitize for Apo2L/TRAIL- or Anticancer Drug-Induced Apoptosis and Induce Regression of Malignant Glioma in Vivo. *Nature Medicine*, 8, 808-815.
<https://doi.org/10.1038/nm735>

[77] Stirpe, F., Olsnes, S. and Pihl, A. (1980) Gelonin, a New Inhibitor of Protein Synthesis, Nontoxic to Intact Cells. Isolation, Characterization, and Preparation of Cytotoxic Complexes with Concanavalin A. *The Journal of Biological Chemistry*, 255, 6947-6953.

[78] Park, YJ, Chang, LC, Liang, JF, et al. (2005) Nontoxic Membrane Translocation Peptide from Protamine, Low Molecular Weight Protamine (LMWP), for Enhanced Intracellular Protein Delivery: In Vitro and In Vivo Study. *THE FASEB Journal*, 19, 1555-1557.

<https://doi.org/10.1096/fj.04-2322fje>

[79] Lu, PY, Xie, FY and Woodle, MC (2003) siRNA-Facilitated Antitumorigenesis for Drug Target Validation and Therapeutics. *Current Opinion in Molecular Therapeutics*, 5, 225-234.

[80] Kim, WJ, Christensen, LV, Jo, S., et al. (2006) Cholesteryl Oligoarginine Delivering Vascular Endothelial Growth Factor siRNA Effectively Inhibits Tumor Growth in Colon Adenocarcinoma. *Molecular Therapy*, 14, 343-350.

<https://doi.org/10.1016/j.ymthe.2006.03.022>

[81] Crombez, L., Morris, MC, Dufort, S., et al. (2009) Targeting Cyclin B1 through Peptide-Based Delivery of siRNA Prevents Tumor Growth. *Nucleic Acids Research*, 37, 4559-4569.

<https://doi.org/10.1093/nar/gkp451>

[82] Josephson, L., Tung, CH, Moore, A., et al. (1999) High-Efficiency Intracellular Magnetic Labeling with Novel Superparamagnetic-Tat Peptide Conjugates. *Bioconjugate Chemistry*, 10, 186-191.

<https://doi.org/10.1021/bc980125h>

[83] Torchilin, VP (2008) Cell

Penetrating Peptide-Modified Pharmaceutical Nanocarriers for Intracellular Drug and Gene Delivery. *Biopolymers*, 90, 604-610.

<https://doi.org/10.1002/bip.20989>

[84] Balzeau, J., Pinier, M., Berges, R., et al. (2013) The Effect of Functionalizing Lipid Nanocapsules with NFL-TBS. 40-63 Peptide on Their Uptake by Glioblastoma Cells. *Biomaterials*, 34, 3381-3389 .

<https://doi.org/10.1016/j.biomaterials.2013.01.068>

[85] Nakase I., Noguchi K., Aoki A., Takatani-Nakase T., Fujii I., Futaki S. (2017). Arginine-rich cell-penetrating peptide-modified extracellular vesicles for active macropinocytosis induction and efficient intracellular delivery. *Sci. Rep.* 7, 1991. 10.1038/s41598-017-02014-6

[86] Tai W., Gao X. (2017). Functional peptides for siRNA delivery. *Adv. Drug Deliv. Rev.* 110-111, 157–

168. 10.1016/j.addr.2016.08.004

[87] Yoo, J., Lee, D., Gujrati, V., Rejinold, N. S., Lekshmi, K. M., Uthaman, S., Jeong, C., Park, I. K., Jon, S., & Kim, Y. C. (2017). Bioreducible branched poly(modified nona-arginine) cell-penetrating peptide as a

novel gene delivery platform. *Journal of controlled release : official journal of the Controlled Release Society*, 246, 142–154.

<https://doi.org/10.1016/j.jconrel.2016.04.040>

[88] Gao W., Yang X., Lin Z., He B., Mei D., Wang D., et al. (2017). The use of electronic-neutral penetrating peptides cyclosporin A to deliver pro-apoptotic peptide: A possibly better choice than positively charged TAT. *J. Controlled Release* 261, 174–186. 10.1016/j.jconrel.2017.06.018

[89] Zhao H., Wu M., Zhu L., Tian Y., Wu M., Li Y., et al. (2018). Cell-penetrating Peptide-modified Targeted Drug-loaded Phase-transformation Lipid Nanoparticles Combined with Low-intensity Focused Ultrasound for Precision Theranostics against Hepatocellular Carcinoma. *Theranostics* 8, 1892–1910. 10.7150/thno.22386

[90] Lo J. H., Hao L., Muzumdar M. D., Raghavan S., Kwon E. J., Pulver E. M., et al. (2018). iRGD-guided Tumor-penetrating Nanocomplexes for Therapeutic siRNA Delivery to Pancreatic Cancer. *Mol. Cancer Ther.* 17, 2377–2388. 10.1158/1535-7163.MCT-17-1090

[91] Liu Y., Song Z., Zheng N., Nagasaka K., Yin L., Cheng J. (2018). Systemic siRNA delivery to tumors by cell-penetrating α -helical polypeptide-based metastable nanoparticles. *Nanoscale* 10, 15339–15349. 10.1039/C8NR03976C

[92] Hu H., Wang J., Wang H., Tan T., Li J., Wang Z., et al. (2018). Cell-penetrating peptide-based nano-vehicles potentiate lymph metastasis targeting and deep penetration for anti-metastasis therapy.

Theranostics 8, 3597–3610.
10.7150/thno.25608

[93] Gao S., Tian B., Han J., Zhang J., Shi Y., Lv Q., et al. (2019). Enhanced transdermal delivery of lornoxicam by nanostructured lipid carrier gels modified with polyarginine peptide for treatment of carrageenan-induced rat paw edema. *Int. J. Nanomed.* Volume 14, 6135–6150. 10.2147/IJN.S205295

[94] Jeong E. J., Choi M., Lee J., Rhim T., Lee K. Y. (2015). The spacer arm length in cell-penetrating peptides influences chitosan/siRNA nanoparticle delivery for pulmonary inflammation treatment. *Nanoscale* 7, 20095–20104. 10.1039/C5NR06903C

[95] Lee Y. Y., Lee S. Y., Park S.-Y., Choi H.-J., Kim E.-G., Han J.-S. (2018). Therapeutic potential of a phospholipase D1 inhibitory peptide fused with a cell-penetrating peptide as a novel anti-asthmatic drug in a Der f 2-induced airway inflammation model. *Exp. Mol. Med.* 50, 1–11. 10.1038/s12276-018-0187-x

[96] Kim J. Y., Ahn J., Kim J., Choi M., Jeon H., Choe K., et al. (2018). Nanoparticle-Assisted Transcutaneous Delivery of a Signal

Transducer and Activator of Transcription 3-Inhibiting Peptide Ameliorates Psoriasis-like Skin Inflammation. *ACS Nano* 12, 6904–6916. 10.1021/acsnano.8b02330

[97] Järver P., Mäger I., Langel Ü. (2010). In vivo biodistribution and efficacy of peptide mediated delivery.

Trends Pharmacol. Sci. 31 (11), 528–535. 10.1016/j.tips.2010.07.006

[98] Jauset T., Beaulieu M.-E. (2019). Bioactive cell penetrating peptides and proteins in cancer: a bright future ahead. *Curr. Opin. Pharmacol.* 47, 133–140. 10.1016/j.coph.2019.03.014

[99] Saar K., Lindgren M., Hansen M., Eiríksdóttir E., Jiang Y., Rosenthal-Aizman K., et al. (2005). Cell-penetrating peptides: A comparative membrane toxicity study. *Analytical Biochemistry* 345 (1), 55–65. 10.1016/j.ab.2005.07.033

[100] Zorko M., Langel U. (2005). Cell-penetrating peptides: mechanism and kinetics of cargo delivery. *Adv.*

Drug Deliv. Rev. 57 (4), 529–545. 10.1016/j.addr.2004.10.010

[101] Moutal A., François-Moutal L., Brittain J. M., Khanna M., Khanna R. (2015). Differential neuro-protective potential of CRMP2 peptide aptamers conjugated to cationic, hy-

drophobic, and amphipathic cell penetrating peptides. *Front. Cell. Neurosci.* 8. 10.3389/fncel.2014.00471

[102] Warrant A., Cardon S., Burlina F., Sagan S. (2017). Membrane crossing and membranotropic activity of cell-penetrating peptides: dangerous liaisons? *Acc. Chem. Res.* 50 (12), 2968–2975. 10.1021/acs.accounts.7b00455

[103] Erazo-Oliveras A., Muthukrishnan N., Baker R., Wang T.-Y., Pellois J.-P. (2012). Improving the Endosomal Escape of Cell-Penetrating Peptides and Their Cargos: Strategies and Challenges. *Pharmaceuticals* 5 (11), 1177–1209. 10.3390/ph5111177

[104] Horton K. L., Stewart K. M., Fonseca S. B., Guo Q., Kelley S. O. (2008). Mitochondria-penetrating peptides. *Chem. Biol.* 15 (4), 375–382. 10.1016/j.chembiol.2008.03.015

[105] Yang, D., Liu, B. & Sha, H. (2025). Advances and prospects of cell-penetrating peptides in tumor immunotherapy. *Sci Rep* 15, 3392. <https://doi.org/10.1038/s41598-025-86130-8>

